

Express Mail #EL447232051US

APPLICATION FOR UNITED STATES

LETTERS PATENT

**MONOCLONAL ANTIBODY FOR ANALYSIS AND CLEARANCE OF
POLYETHYLENE GLYCOL AND POLYETHYLENE GLYCOL-MODIFIED
MOLECULES**

Inventors:

**Steve ROFFLER
Tian-Lu CHENG
Pin-Yi WU**

RELATED APPLICATIONS

This application claims priority from U.S. Provisional Patent Application Serial Number 60/136,522 which was filed on May 28, 1999. The content of the provisional application Serial Number 60/136,522 is hereby incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention generally relates to a novel antibody against polyethylene glycol (PEG) and a process of using this novel antibody to accelerate the clearance of polyethylene glycol conjugates in the human body in cancer therapy as well as a method to quantify the concentration of PEG modified compounds.

2. Description of the Related Art

Recombinant proteins are increasingly being employed for the therapy of a wide variety of diseases (Nemunaitis, 1997; Fareed *et al.*, 1998; Hudson, 1998; Lin, 1998; Harker, 1999; Sandborn and Hanauer, 1999). Many of the proteins in clinical development are modified by the covalent attachment of methoxypoly(ethylene glycol) (PEG) (Menzel *et al.*, 1993; Bassar *et al.*, 1996; Fareed *et al.*, 1998; Goffin *et al.*, 1999; Harker, 1999), a flexible, linear polymer containing repeating $[-OCH_2CH_2-]$ subunits (Delgado *et al.*, 1992). PEG-modified proteins often exhibit prolonged circulation half-lives (Beckman *et al.*, 1988;

Cheng *et al.*, 1997; Chapman *et al.*, 1999) and reduced proteolytic cleavage (Roseng *et al.*, 1992; Kaneda *et al.*, 1995; Brinckerhoff *et al.*, 1999). Immune responses against proteins can also be decreased by covalent attachment of PEG (Abuchowski *et al.*, 1977a). Reduction of immunogenicity can be an important consideration because even recombinant human proteins can induce a humoral immune response (Atkins *et al.*, 1986; Gribben *et al.*, 1990). Conjugates formed between drugs and PEG have also recently been developed (Caliceti *et al.*, 1993; Conover *et al.*, 1997; Greenwald *et al.*, 1998; Pendri *et al.*, 1998). In addition, covalent attachment of PEG to liposomes has been found to reduce non-specific uptake as well as increase liposome stability and half-life (Kirpotin *et al.*, 1997; Cabanes *et al.*, 1998; Meyer *et al.*, 1998).

Clinical development of PEG-modified proteins requires measurement of the pharmacokinetics in animals and patients. Ideally, the concentration of intact PEG-modified protein should be measured. Simple methods to measure intact PEG-protein conjugates, however, are not available. Analysis of PEG-modified proteins has been problematic (Delgado *et al.*, 1992). Sodium dodecylsulfate polyacrylamide gel electrophoresis can be used to measure the relative size of PEG-modified proteins, but the mobility of PEG-modified proteins is slower than the expected molecular weight (Suzuki *et al.*, 1984; Katre *et al.*, 1987). Conjugates can be indirectly measured by first radiolabeling the protein (Kaneda *et al.*, 1995; Cheng *et al.*, 1997; Yabe *et al.*, 1999) or PEG (Mullin *et al.*, 1997), but radioisotopes pose safety concerns and require special handling. Functional assays can be employed to measure the concentration of the protein component of conjugates (Cheng *et al.*, 1997; Esslinger *et al.*, 1997), but no information is provided about the stability of

covalently attached PEG chains. Methods that measure the number of PEG molecules attached to a protein (Habeeb, 1966; Stocks *et al.*, 1986) require that purified conjugate be employed which is difficult to achieve in pharmacokinetic studies. Methods that directly measure the concentration of PEG are relatively insensitive. Colorimetric methods based on complex formation between barium-iodide and PEG require that proteins are first removed and have detection limits of around 1-5 μg PEG (Childs, 1975). A colorimetric method based on partitioning of a chromophore present in aqueous ammonium ferrothiocyanate reagent can be employed for complex protein mixtures but has a detection limit of 1-5 μg PEG (Nag *et al.*, 1996; Nag *et al.*, 1997). High performance liquid chromatography can detect PEG with a detection limit around 1-5 $\mu\text{g}/\text{ml}$ (Kinahan and Smyth, 1991; Ryan *et al.*, 1992; Ruddy and Hadzija, 1994; Miles *et al.*, 1997). Phase-partitioning can be employed to measure PEG but the assay sensitivity is about 1 μg PEG (Guermant *et al.*, 1995). Finally, polyclonal antibodies against PEG can detect the presence of 1 $\mu\text{g}/\text{ml}$ PEG in PEG-modified proteins (Richter and Akerblom, 1983).

The development of monoclonal antibodies against PEG could allow quantitation of PEG and PEG-modified proteins by standard immunological assays including ELISA, immunoblotting, dot blotting and radioimmunoassay. It is difficult, however, to produce antibodies against PEG due to its ability to modulate immune responses. For example, immune responses against proteins are often decreased by covalent attachment of PEG (Abuchowski *et al.*, 1977b). PEG modification has been shown to reduce the immunogenicity of enzymes (Abuchowski *et al.*, 1977a; Chaffee *et al.*, 1992), antibodies (Kitamura *et al.*, 1991), toxins (Wang *et al.*, 1993; He *et al.*, 1999), recombinant human

proteins (Katre, 1990) and other proteins (Chinol *et al.*, 1998). PEG-modified proteins often induce tolerance to the unmodified protein in many (Lee *et al.*, 1981; Savoca *et al.*, 1984; Maiti *et al.*, 1988; Saito *et al.*, 1996; Ito *et al.*, 1997; Ito *et al.*, 1998), but not all cases (Savoca *et al.*, 1979; Chen *et al.*, 1981). Rabbit polyclonal antibodies against PEG have
5 been generated by immunizing rabbits with PEG linked to different proteins (Richter and Akerblom, 1983). PEG alone did not generate an immune response. PEG linked to bovine superoxide dismutase or ragweed pollen extract did not reproducibly generate an anti-PEG response even when the immunogens were given in Freund's complete adjuvant. PEG-modified ovalbumin in Freund's complete adjuvant produced an antibody response in most
10 but not all animals. Antisera raised against PEG-modified ovalbumin could detect PEG-modified proteins with a limit of detection of approximately 1 $\mu\text{g/ml}$ (Richter and Akerblom, 1983). However, no monoclonal antibodies have been successfully produced against PEG to date. Monoclonal antibodies possess advantages compared to polyclonal anti-serum for standardized assays including the ability to produce unlimited quantities of
15 homogeneous antibody with consistent binding affinity.

A major goal of anti-tumor drug development is to increase the therapeutic index of chemotherapy, thereby improving treatment efficacy. One approach to increase the therapeutic index of chemotherapy is to preferentially activate antineoplastic prodrugs at cancer cells but not normal tissues. Tumor selectivity may be achieved by enzymatically
20 converting prodrugs possessing low toxicity to highly toxic anti-neoplastic agents by previously administered antibody-enzyme conjugates (immunoenzymes) that have been allowed to accumulate at tumor cells (Bagshawe *et al.*, 1988; Senter *et al.*, 1988). Even

though maximum accumulation of immunoenzymes in tumors occurs around 24 h after administration (Bosslet *et al.*, 1994; Wallace *et al.*, 1994), prodrugs are generally administered from 3-7 days (Bosslet *et al.*, 1994; Svensson *et al.*, 1998) to up to 2 weeks (Eccles *et al.*, 1994) later to allow adequate time for conjugate to clear from the blood, thereby minimizing systemic prodrug activation and associated toxicity to normal tissues. The requisite of low circulating levels of immunoenzyme therefore often precludes prodrug administration when maximum localization has been achieved.

Prodrugs can be administered during the period of maximum tumor accumulation of immunoenzymes if circulating conjugates are removed or deactivated. Several methods have been devised to accelerate the clearance of radioimmunoconjugates and immunoenzymes from the circulation including the administration of polyclonal (Stewart *et al.*, 1990) and anti-idiotypic antibodies (Ullen *et al.*, 1995a; Ullen *et al.*, 1995b) against the antibody portion of the immunoconjugate, injection of avidin to clear biotinylated antibodies (Klibanov *et al.*, 1988; Paganelli *et al.*, 1991; Stella *et al.*, 1994; Kobayashi *et al.*, 1995), use of monoclonal antibodies against enzymes to clear (Kerr *et al.*, 1993; Haisma *et al.*, 1995) or deactivate (Sharma *et al.*, 1990) immunoenzymes, and extracorporeal immunoabsorption (Tennvall *et al.*, 1997) to remove immunoconjugates from plasma.

β -Glucuronidase targeted to tumor cells can activate the glucuronide prodrug (BHAMG) of p-hydroxy aniline mustard (pHAM) in vitro (Wang *et al.*, 1992) and cure advanced hepatoma ascites in a rat model (Chen *et al.*, 1997). Antibody- β G conjugates injected intravenously, however, are rapidly cleared from the circulation before significant tumor accumulation occurs (Cheng *et al.*, 1997). Modification of β G with PEG can extend

the half-life of antibody- β G conjugates, decrease the normal tissue uptake, and increase the localization of the conjugate at the solid tumors in nude mice (Cheng *et al.*, 1997). Although the extended half-life of PEG-modified β G conjugate is desirable for improved tumor uptake, at least five days were required for the serum concentration of the conjugates to reach a safe level before a prodrug could be administered (Cheng *et al.*, 1997). The accelerated clearance of conjugates from the circulation may allow earlier prodrug administration when the maximum amount of the conjugate is present at tumor cells. We generated mAbs against β G and PEG and examined their effects on the clearance of the β G-sPEG conjugates. The effect of incorporating galactose residues in the clearing antibodies was also examined because galactose can accelerate the removal of proteins from the circulation (Thornburg *et al.*, 1980) by the hepatic asialoglycoprotein receptor (Ong *et al.*, 1991). Rapid clearance of an antibody-carboxypeptidase G2 conjugate by a galactose-modified antibody (Sharma *et al.*, 1990; Sharma *et al.*, 1994) allowed earlier administration of a prodrug with reduced toxicity (Rogers *et al.*, 1995). In addition, galactose-modified mAb has also been employed in a clinical trial to remove residual antibody-carboxypeptidase G2 conjugate from the circulation before prodrug administration (Martin *et al.*, 1997).

SUMMARY OF THE INVENTION

We have produced a monoclonal antibody AGP3 that binds to polyethylene glycol (PEG) by immunizing mice with a conjugate (RH1-BG-PEG) in which PEG-modified beta-glucuronidase (BG) was covalently attached to a murine immunoglobulin (mAb RH1).

5 Fusion of myeloma cells with spleen cells isolated from mice immunized with RH1-BG-PEG without adjuvant resulted in the isolation of a hybridoma secreting an IgM monoclonal antibody to PEG. Such a monoclonal antibody is useful for quantifying the concentration of PEG or PEG-containing compounds in a sample in vitro.

10 In the present application, we have also developed a clearance system for PEG-modified compounds (Cheng *et al.*, 1999; Cheng *et al.*, In press) for use in cancer therapy through antibody-directed enzyme activation of glucuronide prodrugs. We have found that an IgM antibody against PEG can satisfactorily clear PEG-containing compounds from the circulation without significant toxic side-effects.

15 PEG-containing proteins often exhibit extended serum half-lives, reduced immunogenicity and decreased susceptibility to proteolytic degradation (Delgado *et al.*, 1992). Antibody fragments and immunoconjugates that have been modified with PEG also display reduced normal tissue uptake and enhanced tumor accumulation (Pedley *et al.*, 1994; Delgado *et al.*, 1996; Cheng *et al.*, 1997). Clearance of PEG-modified immunoconjugates with AGP3 may therefore be generally useful. We examined whether
20 AGP3 could improve tumor-blood ratios and allow earlier administration of a prodrug in a preclinical model of human colorectal carcinoma.

Accordingly, an object of the present invention is directed to a novel anti-PEG antibody and the preparation thereof.

Another object of the present invention is directed to develop an immunoassay to identify or quantify PEG in a sample, using an anti-PEG monoclonal antibody.

5 A further object of the present invention is directed to a method of treating a cancer patient, which comprises the steps of:

- a. administering a PEG-containing compound comprising tumor targeting means and means for activating an anti-tumor prodrug to the patient;
- 10 b. administering an anti-polyethylene glycol antibody to the patient to accelerate the clearance of the PEG-containing compound from the blood circulation after step a; and
- c. administering the anti-tumor prodrug to said patient after step b.

In a preferred embodiment, PEG-modified BG was covalently linked to the F(ab')₂ fragment of mAb B72.3, an IgG₁ antibody that binds to TAG-72 antigen expressed on the majority of colon adenocarcinomas, invasive ductal carcinomas of the breast, non-small cell lung carcinomas, common epithelial ovarian carcinomas, and gastric, pancreatic and esophageal cancers with limited reactivity to normal adult tissues (Thor *et al.*, 1986). ¹¹¹In-labeled B72.3 (Cyt-103) is approved for clinical use as an imaging agent for colorectal and ovarian carcinoma (Divgi *et al.*, 1994). We have showed that clearance of a PEG-modified B72.3 immunoenzyme with AGP3 produced higher tumor-blood ratios without sacrificing tumor accumulation, allowing earlier prodrug administration with minimal toxicity.

The various features of novelty which characterize the invention are pointed out with particularity in the claims annexed to and forming a part of the disclosure. For a better understanding of the invention, its operating advantages, and specific objects attained by its use, reference should be had to the drawing and descriptive matter in which there are
5 illustrated and described preferred embodiments of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

In the drawings:

Fig. 1 shows binding specificity of antibodies 1E8 and AGP3;

Fig. 2 is an immunoblot showing the binding of antibodies 1E8 and AGP3;

Fig. 3 is an immunoblot of different β G-PEG preparations;

Fig. 4 is an immunoblot showing that AGP3 binds to the backbone of PEG
rather than the linker or methoxy functions groups;

Fig. 5 shows the activity of biotin-labeled AGP3;

Fig. 6 shows measurement of PEG-modified protein by AGP3 ELISA;

Fig. 7 shows measurement of β G-PEG pharmacokinetics by AGP3 ELISA;

Fig. 8 shows the elimination of the galactose-modified antibodies from the
serum of mice;

Fig. 9 shows the clearance of β G-PEG by antibodies;

Fig. 10 shows the dose dependency of the clearance of β G-PEG by AGP3;

Fig. 11 shows the BHAMG-induced organ pathology after β G-PEG clearance;

Fig. 12 shows the weight changes of the mice and hematological toxicity of
BHAMG after the clearance of β G-PEG;

Fig. 13 shows the pH sensitivity of β G;

Fig. 14 shows the activity of B72.3- β G-PEG and H25- β G-PEG.

Fig. 15 shows the specificity of mAb B72.3 to LS174T xenografts;

Fig. 16 shows the sensitivity of LS174T tumor cells to pHAM and BHAMG;

Fig. 17 shows the biodistribution of [¹²⁵I]-labeled B72.3-βG-PEG and H25-βG-PEG in LS174T tumor-bearing BALB/c *nu/nu* mice;

Fig. 18 shows the *in vivo* clearance of the PEG conjugates by the AGP3 antibody;

5 Fig. 19 shows the effect of the clearance of the PEG conjugates by the AGP3 antibody on the tumor localization; and

Fig. 20 shows the *in vivo* antitumor effect of B72.3-βG-PEG conjugate after clearance with AGP3.

DETAILED DESCRIPTION OF THE PRESENTLY PREFERRED EMBODIMENTS

Abbreviations:

1E8-xG, mAb 1E8 containing x galactose moieties per antibody.

5A8-PEG, conjugate formed between mAb 5A8 and β G-sPEG.

5 AGP3-biotin, mAb AGP3 derivatized with biotin.

AGP3-xG, monoclonal antibody AGP3 containing x galactose groups per antibody.

B72.3- β -PEG, conjugate of the F(ab')₂ fragment of mAb B72.3 with β G-sPEG.

β G, *Eschericia coli* β -glucuronidase.

10 β G-PEG- β G, β -glucuronidase crosslinked with poly(ethylene glycol)-di-succinimidyl succinamide.

β G-pPEG, β -glucuronidase modified with pPEG.

β G-sPEG, β -glucuronidase modified with sPEG.

β G-tPEG, β -glucuronidase modified with tPEG.

BMAMG, glucuronide prodrug of pHAM.

15 BSA, bovine serum albumin.

BSA-sPEG, bovine serum albumin modified with sPEG.

ELISA, enzyme-linked immunosorbent assay.

H25- β G-PEG, conjugate of the F(ab')₂ fragment of mAb H25B10 with β G-sPEG.

HRP, horse-radish peroxidase.

20 HRP-GAM, horse-radish peroxidase conjugated goat anti-mouse antibody.

HRP-GAM μ -chain, horse-radish peroxidase conjugated goat anti-mouse IgM μ -chain antibody.

IC₅₀, drug concentration causing 50% inhibition of cellular protein synthesis.

mAb, monoclonal antibody.

PBS, phosphate-buffered saline.

PBS-T, PBS containing 0.05% Tween-20.

5 PEG, monomethoxy poly(ethylene glycol).

PEG-(SSA)₂, poly(ethylene glycol)-di-succinimidyl succinamide

pHAM, *p*-hydroxy aniline mustard.

PNPG, *p*-nitrophenol β -D-glucuronide.

pPEG, methoxypoly(ethylene glycol) succinimidyl propionate.

10 PLT, platelets.

RBC, red blood cells.

RH1- β G-PEG, conjugate formed between mAb RH1 and β G-sPEG.

SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis.

SMCC, succinimidyl-4-[N-maleimidomethyl]cyclohexane-1-carboxylate.

15 sPEG, methoxypoly(ethylene glycol) succinimidyl succinate.

TAG-72, tumor-associated glycoprotein.

tPEG, methoxypoly(ethylene glycol) tresylate.

WBC, white blood cells.

Definitions:

20 As hereinafter used, the term "conjugate" means a complex formed by chemically linking two or more proteins; the term "derivatize" means chemically linking a small

molecular weight compound to a protein; the term "modify" means chemically linking a chemical such as PEG or galactose to the surface of a protein.

Experimental Procedures

Materials: BSA (Fraction V), methoxypoly(ethylene glycol) succinimidyl succinate (sPEG, MW 5000), methoxypoly(ethylene glycol) tresylate (tPEG, MW 5000), cyanomethyl-2, 3, 4, 6-tetra-O-acetyl-1-thio- β -D-galactopyranoside, Ultrogel AcA 22, trinitrobenzenesulfonic acid, fluorescamine, and p-nitrophenyl β -D-glucuronide were purchased from Sigma Chemical Co. (St. Louis, MO). Methoxypoly(ethylene glycol) succinimidyl propionate (pPEG, MW 5000) was obtained from Fluka (Buchs, Switzerland). Poly(ethylene glycol)-di-succinimidyl succinamide (PEG-(SSA)₂, MW 3,4000) was purchased from Shearwater Polymers (Huntsville, AL). Succinimidyl-6-(biotinamido) hexanoate (NHS-LC-biotin) was obtained from Pierce (Rockford, IL). Sephacryl S-200 HR, Sephacryl S-300 HR and Sephadex G-25 gels were from Pharmacia Biotech Asia Pacific Ltd. (Taiwan). pHAM and BHAMG were synthesized as described (Roffler *et al.*, 1991). Succinimidyl-4-[N-maleimidomethyl]cyclohexane-1-carboxylate (SMCC) was from Pierce Chemical Company, Rockford, IL.

Cells: LS174T colon adenocarcinoma cells (ATCC CL-188) were obtained from the American Type Culture Collection (Rockville, MD). Cells were mycoplasma free and cultured in Dulbecco's modified Eagle's medium (Gibco BRL, Grand Island, NY) supplemented with 10% heat-inactivated bovine serum, 100 U/ml penicillin and 100 μ g/ml streptomycin. FO myeloma cells (ATCC CRL-1646) were purchased from the American Type Culture Collection (Manassas, VA).

Animals: BALB/c mice were obtained from the animal room of the Institute of Biomedical Sciences, Academia Sinica. BALB/c *nu/nu* mice were from the Cancer

Research Laboratory, Tri-Service General Hospital, Taipei. Animal experiments were performed in accordance with institute guidelines.

Apparent to those ordinarily skilled in the art, any of the above described materials, cell lines or animals may be substituted by other like materials, cell lines, animals or their equivalents, produced by any manufacturer, to perform substantially the same functions.

Antibodies: mAb RH1, an IgG_{2a} mAb that binds to an antigen expressed on the surface of AS-30D rat hepatoma cells, was generated as described (Roffler *et al.*, 1994). Hybridomas secreting mAb B72.3 (ATCC HB-8108), an IgG₁ monoclonal antibody specific to the tumor-associated glycoprotein (TAG-72) antigen (Johnson *et al.*, 1986) and mAb H25B10 (ATCC CRL-8017), a control IgG₁ antibody that binds to the surface antigen of hepatitis B virus, were obtained from the American Type Culture Collection. mAb 5A8, an IgG_{2a} mAb that binds to the surface immunoglobulin expressed on 38C13 lymphoma cells (Maloney *et al.*, 1985), was provided by Dr. Mi-Hua Tao, Institute of Biomedical Sciences, Academia Sinica. mAbs were purified from ascites by affinity chromatography on Sepharose-protein A in high salt buffer. The F(ab')₂ fragments of mAb B72.3 and H25B10 were generated by proteolytic digestion of whole antibodies with bromelain (Milenic *et al.*, 1989). F(ab')₂ fragments were purified by sequential chromatography on DE-52 (anion exchange) and Sephacryl S-200 HR (gel filtration). Streptavidin-HRP (horse radish peroxidase) was from Serotec (Oxford, UK). HRP-conjugated goat anti-mouse IgM μ -chain antibody (HRP-GAM μ -chain) and HRP-conjugated goat anti-mouse IgG (HRP-GAM) were from Organon Teknika (Durham, NC).

PEG-Modification of β G: Recombinant *Eschericia coli*-derived β G was produced and purified as described (Cheng *et al.*, 1997). Twenty milligrams of sPEG per milligram of β G (2 mg/mL) were mixed in coupling buffer (deoxygenated PBS containing 1 mM EDTA, pH 8.0) at room temperature for 2 h before one-tenth the volume of a saturated glycine solution was added to stop the reaction. Unreacted PEG was removed by dialysis against PBS or by gel filtration on a 2.5 x 100 cm Sephacryl S-200 HR column eluted with PBS at 15 mL/h. Relevant fractions were pooled and concentrated by ultrafiltration to 1.5-2.0 mg/mL for storage at -80 °C. BSA was modified with sPEG in a similar manner. In some experiments, β G in coupling buffer was modified with 2 mg of sPEG or with 2 or 20 mg of tPEG/mg of β G.

To produce β G-pPEG, β G was passed through a 2.5 x 30 cm Sephadex G-25 column equilibrated with 0.1 M borate buffer, pH 8.0 and concentrated by ultrafiltration to 3 mg/ml. Ten milligrams pPEG in 0.1 M borate buffer, pH 8.0 (150 mg/ml) was added per mg β G and mixed for 2 h at room temperature before one-tenth volume of a saturated solution of glycine was added to stop the reaction. Unreacted PEG was removed by gel filtration on a 2.5 x 100 cm Ultrogel ACA 22 column eluted with PBS at 15 ml/h. Relevant fractions containing β G-PEG were pooled and concentrated by ultrafiltration to 1 mg/ml for storage at -80 °C. β G was crosslinked with PEG-(SSA)₂ (3.3 mg PEG per mg β G) in a similar manner to produce β G-PEG- β G.

Protein concentrations were determined by the bicinchoninic acid assay (Pierce, Rockford, IL) with BSA employed as the reference protein. The BCA assay was not affected by covalently coupled PEG after removal of free PEG.

Conjugation of antibodies to β G-sPEG: mAb RH1 and 5A8 were passed through a 2.6 x 30 cm Sephadex G-25 column equilibrated with coupling buffer and concentrated by ultrafiltration to 3 mg/mL. A 4-fold molar excess of SMCC (1 mg/mL in dioxane) was slowly added to the antibodies and allowed to react at room temperature for 50 min.

5 Unreacted SMCC was removed by gel filtration on a 2.6 x 30 cm Sephadex G-25 column equilibrated with coupling buffer. Typically, an average of 1 to 1.5 maleimido groups was introduced into antibodies. β G-sPEG was partially reduced by adding dithiothreitol to a final concentration of 20 mM for 1 h at 37 °C. Freshly derivatized antibody and β G-sPEG were immediately mixed at an equal molar ratio, concentrated to 2 mg/mL by ultrafiltration
10 and incubated at room temperature for 2 h. Cysteine was added to a final concentration of 2 mM to stop the reaction and the mixture was concentrated by ultrafiltration to about 5 mg/mL. Conjugates were purified on a 2.5 x 100 cm Sephacryl S-300 HR column equilibrated with PBS at a flow rate of 15 mL/h. Fractions containing conjugate monomers (antibody/enzyme = 1:1) were pooled and concentrated by ultrafiltration to 1.5-2.0
15 mg/mL for storage at -80 °C. The yield of conjugates averaged around 35% based on the weight of the purified conjugate divided by the total starting weights of antibody and enzyme.

mAb B72.3-F(ab')₂ and mAb H25-F(ab')₂ were also linked to β G-sPEG by the procedure described above to produce the conjugates B72.3- β G-PEG and H25- β G-PEG.

20 Conjugates were purified by size exclusion chromatography on Sephacryl S-300 HR equilibrated with PBS. The antigen-binding activity of conjugates were measured by ELISA in 96-well microtiter plates coated with bovine submaxillary gland mucin (King *et*

al., 1994). The combined antigen-binding and enzymatic activities were measured as described (Chen *et al.*, 1997).

Generation of Monoclonal Antibodies against β G and β G-PEG: Generally, monoclonal antibodies against a PEG molecule or moiety may be produced by conjugating the PEG molecule with an antigenic molecule, such as a protein. Immunization of mice with such conjugated PEG to produce desired antibodies can be routinely performed by a person of ordinary skill in the art. Preferably, the conjugated PEG is linked with another molecule to form a three-part complex for immunization of mice. Most preferably, female BALB/c mice were i.v. injected with 200 μ g of RH1- β G-PEG in PBS on day 1 and i.p. injected with 100 μ g of RH1- β G-PEG on days 7, 14, 21 and 28. Mice received a final i.v. injection of 200 μ g of RH1- β G-PEG on day 54. Hybridomas were generated by fusing spleen cells with FO myeloma cells 3 days later (Yeh *et al.*, 1979). Hybridomas were screened by ELISA in 96-well microtiter plates coated with β G or β G-PEG. Positive hybridomas were cloned by limiting dilution in 96-well plates containing thymocyte feeder cells in HT medium supplemented with 15% fetal calf serum. The isotypes and subclasses of mAbs were determined with the Mouse MonoAb-ID kit according to the manufacturer's instructions (Zymed laboratories, South San Francisco, CA). The apparent affinities of antibodies were measured as described (Beatty *et al.*, 1987).

Purification of AGP3: AGP3 ascites obtained from pristane-primed eight-week old BALB/c female mice was centrifuged at 5000 g for 10 min to remove debris. Ascites fluid (5 ml) was passed through a 5 ml Sephadex G-25 column equilibrated with PBS and then purified by gel filtration on a 2.5 x 100 cm Ultrogel AcA 22 column eluted with PBS at 15

ml/h. Fractions containing AGP3 were collected, concentrated by ultrafiltration to 1-2 mg/ml and stored at -80 °C. Yields of 5-6 mg AGP3 per ml ascites were typically achieved.

Biotinylation of AGP3: AGP3 (1 mg/ml) in 0.1 M borate buffer, pH 8.0 was mixed
5 with a 25-fold molar excess of NHS-LC-biotin for 1 h at room temperature to produce AGP3-biotin. One-tenth volume of a saturated solution of glycine was added to stop the reaction before aliquots were stored at -80 °C. The antigen-binding activity of AGP3-biotin was determined by ELISA in 96-well microplates coated overnight with 1 µg/well
10 βG-tPEG (20 mg tPEG per mg βG). Plates were blocked 1 h at 37 °C with 2% skim milk in PBS before 50 µL/well samples of AGP3 or AGP3-biotin in dilution buffer (0.5% BSA, 0.05% Tween-20 in PBS) were added to the wells for 1 h at room temperature. Plates were washed 6 times with PBS/TB (0.05% bovine serum albumin, 0.1% Tween-20 in PBS) and 50 µL/well of either streptavidin-HRP (1:1000) or HRP-GAM µ-chain (1:1000) in dilution
15 buffer were added for 1 h at room temperature. The plates were washed 6 times and 100 µL/well ABTS substrate (0.4 mg/ml 2,2'-azino-di(3-ethylbenzthiazoline-6-sulfonic acid), 0.003% H₂O₂, 100 mM phosphate-citrate, pH 4.0) was added for 30 min at room temperature. Absorbance (405 nm) of wells was measured in a Molecular Devices (Menlo Park, CA) microtiter plate reader.

Radiolabeling of AGP3: AGP3 may be radiolabeled by a radioactive isotope such as
20 ¹²⁵I or ¹³¹I. Any established method such as a method of iodination suitable for radiolabeling a monoclonal antibody may be used for labeling AGP3.

AGP3 ELISA: MaxisorpTM 96-well microplates (NuncTM, Roskilde, Denmark)

were coated with 50 μ L/well of AGP3 (200 μ g/ml in PBS) for 3 h at 37 °C. Plates were blocked with 2% skim milk in PBS overnight and washed twice with PBS/T (0.05% Tween-20 in PBS) before serial dilutions of BG or BG-pPEG (50 μ L in dilution buffer) were added to wells for 1.5 h at room temperature. Plates were washed 6 times with PBS/T and 50 μ L/well AGP3-biotin (20 μ g/ml in dilution buffer) was added for 1.5 h at room temperature. Plates were again washed with PBS/T and 50 μ L/well streptavidin-HRP (1:1000) was added for 1 h at room temperature. Plates were washed before 100 μ L/well ABTS substrate was added for 30 min at room temperature. Absorbance (405 nm) of wells was measured in a microtiter plate reader. The effect of serum on the AGP3 ELISA was examined by performing the assay in the presence of 10% serum isolated from healthy BALB/c mice.

The affinities of AGP3 to various PEG molecules were examined using ELISA. Generally, AGP3 may bind any PEG molecule with a tendency that its affinities to those PEG molecules increase with PEG's molecular weights, i.e., the higher the PEG molecular weight, the better AGP3 will bind. Preferably, AGP3 binds PEG molecules having molecular weights ranging from 3,000 to 60,000 daltons, and molecular weights ranging from 5,000 to 60,000 daltons are most preferred.

Pharmacokinetics of BG-pPEG: Male BALB/c mice were i.v injected with 150 μ g BG-pPEG. Serum was isolated from blood samples that were periodically removed from the tail vein of the mice. The concentration of BG-pPEG in serially-diluted serum samples was assayed in triplicate by the AGP3 ELISA. The concentration of BG-pPEG was

determined by comparison to a standard curve generated with known concentrations of β G-pPEG.

Immunoblotting of PEG-Modified Proteins: Samples were electrophoresed in sodium-dodecyl polyacrylamide gels under reducing conditions before overnight transfer to nitrocellulose paper by passive diffusion in blotting buffer (50 mM NaCl, 2 mM EDTA, 0.5 mM 2-mercapto-ethanol, and 10 mM Tris-HCl, pH 7.5). Blots were blocked with 5% skim milk and incubated for 1 h at 37 °C with 1E8 or AGP3 ascites diluted 1:2000 in PBS containing 0.05% BSA. Blots were washed 3 times with PBS-T and twice with PBS before incubation with HRP-GAM or HRP-GAM μ -chain for 30 min at 37 °C. Blots were washed and bands were visualized by ECL detection according to the manufacturer's instructions (Pierce, Rockford, IL).

Galactose-Modification of Antibodies: Cyano-methyl-2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-galactopyranoside was reacted with 0.1 M sodium methoxide in dry methanol for 48 h (Marshall *et al.*, 1995). The methanol was removed by evaporation under vacuum and 1E8 or AGP3 (5 mg/mL in pH 8.5 borate buffer) were directly added and allowed to react for 2 h. Different amounts of galactose were incorporated into antibodies by varying the ratio of the reactants. The number of galactose groups introduced into antibodies was estimated by the reduction of trinitrobenzenesulfonic acid reactive amine groups after galactose modification (Habeeb, 1966) using a molar extinction coefficient (ϵ^{335}) of 1.31×10^4 for 1E8 and 1.38×10^4 for AGP3. Galactose groups were also estimated by measuring amine groups with flouorescamine before and after galactose modification (Stocks *et al.*, 1986).

Serum Half-Lives of Galactose-Modified 1E8 and AGP3: BALB/c mice were i.v.

injected with 200 μ g. of unmodified or galactose-modified 1E8 or AGP3. Serum was isolated from blood samples that were periodically removed from the tail vein of mice. The concentrations of 1E8 or AGP3 were determined in duplicate samples by ELISA in microtiter plates coated with β G or β G-sPEG, respectively, by comparison to standard curves generated with known concentrations of the corresponding unmodified or galactose-modified antibodies.

Clearance of β G-PEG and 5A8- β G-PEG from Serum: Groups of 3 BALB/c mice were i.v. injected with β G-sPEG or 5A8- β G-PEG at time zero. Blood samples were periodically removed before i.v. injection of unmodified or galactose-modified 1E8 or AGP3 at 24 h. Additional blood samples were taken at subsequent times and the β G activity in duplicate samples was measured using *p*-nitrophenol β -D-glucuronide as substrate (Wang *et al.*, 1992). Sample concentrations were calculated by comparison of absorbance values with standard curves constructed from known concentrations of β G-sPEG or 5A8- β G-sPEG. For meaningful comparison of clearance results, the measured concentrations of β G-sPEG in mice injected with clearing antibodies were normalized to the mean concentration of β G-sPEG in control (uncleared) mice just before clearance by

$$N_j = (C_{mi}/C_i)C_j$$

where N_j is the normalized serum concentration of β G-sPEG in an experimental mouse at time j , C_{mi} is the mean serum concentration of β G-sPEG in the control group serum at the last time point before clearance, C_i is the serum concentration of β G-sPEG in an

experimental mouse at the last time point before clearance, and C_j is the serum concentration of BG-sPEG in an experimental mouse at time j .

Toxicity of Clearance: The numbers of WBC, RBC, and PLT in groups of 6 BALB/c mice were measured as described (Chen *et al.*, 1997) 1 day before i.v. injection of 50 μ g of BG-PEG or PBS. Mice were i.v. injected 24 h later with 300 μ g AGP3 or AGP3-208G, or 2 fractionated 150 μ g doses of AGP3-208G separated by 4 h. After allowing 6 h for clearance, mice were i.v. injected with two 7.5 mg/kg doses of BHAMG in PBS. On day 8, the numbers of WBC, RBC, and PLT in serum samples were determined. On days 3 and 10, the spleen, liver and kidney were removed from one mouse in each group and fixed in 10% formalin. Ethanol dehydrated samples were embedded in paraffin and cut into 3 μ m sections. Sections were then dewaxed with xylene, stained with hematoxylin/eosin, and dehydrated before observation under a light microscope. The weight of mice was followed for 18 days.

pH Dependence of BG Activity: Fifteen μ L of BG (4 μ g/mL) was mixed in microtiter plates with 200 μ l of BG buffer (Wang *et al.*, 1992) with pH values ranging from 3-10. Twenty-five μ L of *p*-nitrophenol β -D-glucuronide (32 mM) was added to the wells at 37 $^{\circ}$ C for 15 minutes, excess 1N NaOH was added to adjust the pH to greater than 10, and absorbance was read at 405 nm in a microplate reader. To test the pH stability of BG, enzyme in BG buffer was adjusted to pH values ranging from 3-11 for 4 h at 37 $^{\circ}$ C before neutralization to pH 7.0 and assay of enzymatic activity.

Drug sensitivity of LS174T cells: LS174T cells were plated overnight in 96 well microtiter plates at 40,000 cells per well. Serial dilutions of pHAM or BHAMG in medium

containing 10% fetal calf serum were added to cells in triplicate for 24 h at 37°C. Cells were subsequently washed once with sterile PBS, incubated until hour 48 in fresh leucine-free medium and then pulsed for 12 h with [³H]-leucine (1μCi/well) in fresh medium and harvested with a Filter-mate (Packard). Incorporated radioactivity was determined on a Top-Count scintillation counter (Packard). Results are expressed as percentage inhibition of [³H]-leucine incorporation compared with untreated cells (control) by the following formula:

$$\% \text{ inhibition} = 100 \times \frac{\text{cpm sample} - \text{cpm background}}{\text{cpm control} - \text{cpm background}}$$

Radiolabeling of Conjugates: B72.3-βG-PEG and H25-βG-PEG were labeled with ¹²⁵I Bolton-Hunter reagent to specific activities of 0.5-0.7 and 0.5-0.7 μCi/μg, respectively, according to the manufacturer's instructions (ICN Pharmaceuticals Inc., Costa Mesa, CA). Conjugates retained antigen-binding activity and specificity as determined by radioimmunoassay against bovine submaxillary gland mucin coated in 96-well microtiter plates. βG activity was unaffected and conjugates were not degraded as determined by gel autoradiography after sodium dodecyl sulfate polyacrylamide electrophoresis.

Tumor Localization of B72.3-βG-PEG: 5 x 10⁶ LS174T cells were injected s.c. in 6-8 week old nude mice. After tumors had grown to approximately 100-200 mm³, 200 μg (60 μCi) [¹²⁵I]B72.3-βG-PEG or [¹²⁵I]H25-βG-PEG were i.v. injected into the lateral tail vein of mice. Groups of 3-4 mice were sacrificed after 24, 48, 72 and 96 h. Tumors, blood and organs were weighed on an analytical balance and assayed for radioactivity in a multichannel gamma-counter. The results are expressed as the specific uptake of the conjugates by the tumor or the tissue (% injected dose/g).

In vivo Clearance of B72.3- β G-PEG and H25- β G-PEG: Groups of 4 BALB/c *nu/nu* mice were i.v. injected with 250 μ g B72.3- β G-PEG or H25- β G-PEG at time zero. Blood samples were periodically removed before mice were given two i.v. injection of 300 and 200 μ g of AGP3 at 48 and 50 h. Additional blood samples were taken at subsequent times and the β G activity in duplicate samples was measured using p-nitrophenol β -D-glucuronide as substrate (Wang *et al.*, 1992). Sample concentrations were calculated by comparison of absorbance values with a standard curve constructed from known concentrations of B72.3- β G-PEG and H25- β G-PEG.

Tumor Localization of B72.3- β G-PEG after Clearance: 5 x 10⁶ LS174T cells were injected s.c. in 6-8 week old nude mice. After 12 days, groups of 8 BALB/c *nu/nu* mice bearing 100-200 mm³ LS174T tumor xenografts were i.v. injected at time 0 with 200 μ g (140 μ Ci) [¹²⁵I] B72.3- β G-PEG or [¹²⁵I] H25- β G-PEG. Half of the mice were i.v. injected 48 and 50 h later with two fractionated 300 and 200 μ g doses of AGP3 whereas PBS was given to the other half. Mice were sacrificed after allowing 6 h for clearance. Tumors, blood and organs were weighed on an analytical balance and assayed for radioactivity in a multichannel γ -counter. Results are expressed as specific uptake of conjugate in tumor or tissues (% injected dose/g).

Therapy of LS174T Tumor Xenografts: Groups of 9 BALB/c *nu/nu* mice were s.c. injected on the right flank with 5 x 10⁶ LS174T cells on day 1. On day 10, mice were i.v. injected via the lateral tail vein with PBS, 250 μ g B72.3- β G-PEG or 250 μ g H25- β G-PEG. 48 and 50 h later mice were i.v. injected with 2 sequential doses of 300 and 200 μ g AGP3. Mice were i.v. injected 6 h later with BHAMG (7.5 mg/kg x 3). Control groups of tumor-

bearing mice were treated with BHAMG (7.5 mg/kg x 3), pHAM (2 mg/kg x 3) or PBS alone. Therapy was repeated starting on days 16, 26, and 41. All mice received 2-4 rounds of therapy. Tumor volumes (length x width x height x 0.5) were measured twice a week and mice weight was followed as an estimate of treatment toxicity. Mice were killed when they displayed signs of morbidity or when the tumor size exceeded 2.5 cm³. Groups of 8 BALB/c *nu/nu* mice bearing larger 200-250 mm³ tumors were also treated as above with 2 rounds of therapy starting on days 11 and 23.

Data Analysis: Statistical significance of differences between mean values was calculated with the shareware program Schoolstat (White Ant Occasional Publishing, West Melbourne, Australia) using the independent *t*-test for unequal variances.

The methods described above are illustrative. They shall not be construed as limitations to the present invention. The present invention encompasses any variations or modifications of the above described methods, which are known or would have been known to those ordinarily skilled in the art.

RESULTS

Antibodies against β G-PEG: Monoclonal antibodies against β G-sPEG were generated from spleen cells of mice that had received multiple injections of RH1- β G-PEG. mAb 1E8 (IgG₁) and AGP3 (IgM) were selected for further characterization based on preliminary screening by ELISA. The binding specificities of 1E8 and AGP3 are shown in Figures 1A and 1B. In Figure 1A, 1E8 (squares) and AGP3 (circles) antibodies were incubated in microtiter plates coated with β G (open symbols) or β G-sPEG (filled symbols). In Figure 1B, 1E8 (squares) and AGP3 (circles) were incubated in microtiter plates coated

with BSA (open symbols) or BSA-sPEG (filled symbols). Binding of antibodies was detected by measuring the absorbance of the wells at 405 nm as described in the Experimental Procedures. The mean absorbance values of triplicate determinations are shown. Bars show the standard error of the mean. 1E8 bound to β G and, to a lesser extent, β G-sPEG (Figure 1A). The apparent affinity constants of 1E8 for β G and β G-sPEG were 0.83 and 1.8 nM, respectively (data not shown). In contrast, AGP3 bound β G-sPEG with an apparent affinity constant of 0.79 nM but did not bind β G (Figure 1A), indicating that this antibody recognizes PEG chains present on β G-sPEG. AGP3 specificity for PEG was supported by the finding that it also bound BSA-sPEG but not BSA (Figure 1B). As expected, 1E8 bound neither BSA nor BSA-sPEG (Figure 1B). Neither of the mAbs affected the activity of β G (results not shown).

Specificity of the mAb binding was further examined by immunoblotting, as shown in Figures 2A and 2B. Proteins were electrophoresed in a 3-12.5% gradient reduced SDS-PAGE, transferred to nitrocellulose paper, and probed with 1E8 (2A) or AGP3 (2B) as described in the Experimental Procedures. Lane 1 is β G; lane 2 is β G-sPEG (2 mg of sPEG/mg of β G); lane 3 is β G-sPEG (20 mg of sPEG/mg of β G); lane 4 is β G-tPEG (2 mg of tPEG/mg of β G); lane 5 is β G-tPEG (20 mg of tPEG/mg of β G); lane 6 is BSA and lane 7 is BSA-sPEG. kDa represents molecular mass in thousands. The results show that 1E8 detected β G and PEG-modified β G (lanes 1-5), but not BSA or BSA-sPEG (lanes 6 and 7). Figure 2B shows that AGP3, in contrast, did not bind β G (lane 1) or BSA (lane 6) but detected all proteins that incorporated PEG (β G-sPEG, β G-tPEG, and BSA-sPEG). The

ability of AGP3 to bind either sPEG or tPEG-modified proteins demonstrates that this antibody binds to poly(ethylene glycol) rather than the linker between PEG and proteins.

Figures 3A and 3B demonstrate immunoblots of different β G-PEG preparations. Samples were electrophoresed on an 8% reduced SDS-PAGE, transferred to nitrocellulose paper, and probed with 1E8 (A) or AGP3 (B) as described in the Experimental Procedures. Lane 1 is β G; lane 2 is β G-sPEG purified by gel filtration; lane 3 is β G-sPEG purified by dialysis; and lane 4 is sPEG neutralized with excess glycine. kDa is molecular mass in thousands. Substantial amounts of free PEG were present in β G-sPEG preparations purified by dialysis (Figure 3B, lane 3). AGP3 immunoblotting of sPEG after neutralization with glycine (Figure 3B, lane 4) produced a similar smeared band, indicating that the low molecular weight material in β G-sPEG corresponded to free PEG rather than to degraded β G-sPEG fragments. Purification of β G-sPEG by gel filtration on Sephacryl S-200 HR resulted in the removal of free PEG (Figure 3B, lane 2). This β G-PEG conjugate, containing 9.2 ± 0.3 sPEG groups/ β G subunit and possessing 69% of original enzymatic activity, was employed in all subsequent experiments using β G-sPEG.

The antigen-binding specificity of AGP3 was further examined by immunoblotting, as shown in Figures 4A and 4B. Proteins were electrophoresed in a 3-12.5% polyacrylamide gel under reducing conditions, transferred to nitrocellulose paper and probed with mAb 1E8 (Fig. 4A) or AGP3 (Fig. 4B) as described in the Experimental Procedures. Lane 1 is β G-PEG- β G (β G modified with polyethylene glycol di-succinimidyl succinamide); lane 2 is unmodified β G; and lane 3 β G-pPEG (β G modified with methoxypolyethylene glycol succinimidyl propionate). The electrophoretic mobility of β G-

pPEG (Fig. 4A, lane 3) was slower than unmodified β G (Fig. 4A, lane 2), demonstrating successful conjugation of pPEG to β G. β G was also successfully crosslinked with PEG to form β G-PEG- β G (Fig. 4A, lane 1). β G-PEG- β G does not contain a terminal methoxy group on the PEG chain. AGP3 bound β G-pPEG (Fig. 4B, lane 3) and β G-PEG- β G (Fig. 4B, lane 1) but not unmodified β G (Fig. 4B, lane 2). Binding of AGP3 to β G-PEG- β G shows that AGP3 binds to the backbone of PEG rather than the methoxy group present at the terminal of mPEG. This finding indicates that multiple AGP3 mAbs can bind to a single PEG chain and prompted us to examine whether AGP3 could be employed in a sandwich ELISA for the detection of PEG-modified proteins.

AGP3 ELISA: AGP3 was examined for use in the ELISA detection of PEG-modified proteins. AGP3 was modified with a 25-fold molar excess of NHS-LC-biotin to produce AGP3-biotin. In Figure 5, AGP3 (squares) or AGP3-biotin (circles) were incubated in a microtiter plate coated with β G-tPEG. Antibody binding to immobilized PEG was detected with HRP-GAM μ -chain antibody (open symbols) or streptavidin-HRP (filled symbols). Results represent average values of duplicate determinations. AGP3-biotin retained the same antigen binding activity to β G-tPEG as AGP3 in ELISA using HRP-GAM μ -chain antibody for detection (Fig. 5). Detection of antibody binding to β G-PEG with streptavidin-HRP showed that AGP3-biotin but not AGP3 could be detected (Fig. 5), demonstrating that biotin was indeed incorporated into AGP3-biotin.

A sandwich ELISA was developed in which AGP3 was employed for the capture step and AGP3-biotin used for detection. In Figure 6, serial dilutions of β G-pPEG in PBS (O) or 10% mouse serum (\square) as well as β G in PBS (Δ) were incubated in a microtiter plate

coated with AGP3. Binding of β G-pPEG to the immobilized AGP3 was detected by sequential addition of AGP3-biotin and streptavidin-HRP. Results represent mean values of triplicate determinations. Bars indicate the standard error of the mean. Figure 6 shows that this assay could detect β G-pPEG at concentrations at least as low as 15 ng/ml. This corresponds to 2.6 fmol of β G-pPEG or 500 pg of PEG. The assay only detected PEG-modified protein as shown by the lack of signal when unmodified β G was assayed (Fig. 6). The effect of serum on the ELISA was examined by adding 10% mouse serum to samples before assay. The sensitivity of the assay was about the same as without serum (Fig. 6), indicating that this ELISA should be applicable to pharmacokinetic studies of PEG-modified molecules and proteins. The AGP3 ELISA was employed to measure the pharmacokinetics of β G-pPEG in mice. In Figure 7, a BALB/c mouse was i.v. injected with 150 μ g β G-pPEG. Blood samples were periodically removed from the tail vein and assayed for β G-pPEG concentration by ELISA. Figure 7 shows that after a period of rapid clearance, β G-pPEG was eliminated from the circulation with a first-order half-life of 19.2 h. This result indicates that the newly developed ELISA can be employed to determine the pharmacokinetics of PEG-modified compounds.

There are currently no methods to directly measure low concentrations of intact PEG-conjugates for pharmacological studies. Conjugates can be indirectly measured by first radiolabeling the protein (Kaneda *et al.*, 1995; Cheng *et al.*, 1997; Yabe *et al.*, 1999) or PEG (Mullin *et al.*, 1997), but radioisotopes pose safety concerns and require special handling. Functional assays can be employed to measure the concentration of the protein component of conjugates (Esslinger *et al.*, 1997; Cheng *et al.*, 1999), but no information is

provided about the stability of covalently attached PEG chains. Methods that measure the number of PEG molecules attached to a protein (Habeeb, 1966; Stocks *et al.*, 1986) require that purified conjugate be employed which is difficult to achieve in pharmacokinetic studies. Methods that directly measure the concentration of PEG are relatively insensitive.

5 Colorimetric methods based on complex formation between barium-iodide and PEG require that proteins are first removed and have detection limits of around 1-5 μg PEG (Childs, 1975). A colorimetric method based on partitioning of a chromophore present in aqueous ammonium ferrothiocyanate reagent can be employed for complex protein mixtures but has a detection limit of 1-5 μg PEG (Nag *et al.*, 1996; Nag *et al.*, 1997). High performance

10 liquid chromatography can measure PEG with a detection limit around 1-5 $\mu\text{g}/\text{ml}$ (Kinahan and Smyth, 1991; Ryan *et al.*, 1992; Ruddy and Hadzija, 1994; Miles *et al.*, 1997). Phase-partitioning can be employed to measure PEG but the assay sensitivity is about 1 μg PEG (Guermant *et al.*, 1995). Finally, polyclonal antibodies against PEG can detect the presence of 1 $\mu\text{g}/\text{ml}$ PEG in PEG-modified proteins (Richter and Akerblom, 1983). The newly

15 developed ELISA can measure 500 pg of PEG in $\beta\text{G-pPEG}$, approximately 4 orders of magnitude more sensitive than other methods. In addition, the AGP3 ELISA does not require prior separation of PEG-conjugates from complex mixtures, simplifying application to pharmacokinetic studies.

Glactose-Modification of Antibodies: 1E8 and AGP3 were derivatized with galactose

20 to target their uptake by the asialoglycoprotein receptor on hepatocytes. Modification of 1E8 with increasing amounts of activated galactose resulted in the incorporation of up to 47 galactose residues/antibody, as shown in Table 1.

Table 1

Characteristics of Galactose-Modified Monoclonal Antibodies

Galatose added ^a (mg/mg)	Conjugate Name	Incorporated galactose		Activity ^d (% control)	Conjugate Name	Incorporated galactose		Activity ^d (% control)
		<u>Groups/mAb</u>				<u>Groups/mAb</u>		
		Fluor ^b	TNBS ^c			Fluor ^b	TNBS ^c	
0	1E8	0	0	100	AGP3	0	0	100
0.9	1E8-9G	12±0		100	AGP3-64G	60±18	67±10	72
1.8	1E8-19G	7±0.6		82	AGP3-130G	130±4	130±13	64
5.5	1E8-36G	17±2		72	AGP3-208G	210±8	205±21	48
10.9	1E8-46G	21±2		62	AGP3-285G	290±0.1	280±8	43
16.4	1E8-47G	35±0.7		66	AGP3-295G	310±0.1	280±7	42
		37±2						
		48±0.1						
		44±3						
		47±0.3						
		46±2						

^a The amount of activated galactose added per milligram of antibody. ^b Determined by the fluorescamine method as described by Stocks

et al. (1986). ^c Determined by the trinitrobenzenesulfonic acid method as described in the Experimental Procedures. ^d Antigen-binding

activity determined by ELISA.

Introduction of up to 20 galactose residues did not affect the binding of 1E8 to β G whereas greater incorporation of galactose decreased 1E8 activity by about 35%. A maximum of about 290 galactose residues could be introduced into AGP3 (Table 1). The greater number of galactose residues incorporated into AGP3 primarily reflects the larger size of the IgM molecule; galactose incorporation for both 1E8 and AGP3 corresponded to 0.3 galactose moieties/kDa of antibody. AGP3 was more sensitive to galactose modification with a 2.4-fold decrease in antigen-binding activity at the highest levels of galactose incorporation.

The half-lives of galactose-modified antibodies were determined in BALB/c mice, as shown in Figures 8A and 8B. Each of the BALB/c mice was i.v. injected with 200 μ g of 1E8 (●), 1E8-9G (□), 1E8-19G (◇), 1E8-36G (Δ), or 1E8-46 (O) (Fig. 8A) or 200 μ g of AGP3 (●), AGP3-64G (□), AGP3-130G (◇), AGP3-208G (Δ), or AGP3-285G (O) (Fig. 8B). The concentration of antibodies in serum samples taken at the indicated times was measured as described in the Experimental Procedures. Figure 8A shows that 1E8 was eliminated from serum with a half-life of 22 h. Introduction of increasing numbers of galactose residues into 1E8 resulted in progressively faster initial clearance rates. Even at the highest galactose incorporation, however, residual concentrations of 1-4 μ g/mL 1E8-46G were present in the serum after 5 h. Unmodified AGP3 cleared from serum with a half-life of 13 h but in contrast to galactose-modified 1E8, almost no residual antibody could be detected in the circulation within 5 h after administration of AGP3 modified with more than 130 galactose moieties (Figure 8B). Increasing the level of galactose residues per antibody

did not greatly affect the initial rate of clearance, but did increase the total amount of antibody that was removed from the circulation. For example, the concentrations of 1E8-9G, 1E8-19G and 1E8-36G in serum were 45%, 5.3% and 1.2% of 1E8 levels after 8 h. The galactose-modified antibodies that were not removed from the circulation within the first 5 h exhibited similar elimination kinetics as unmodified 1E8, indicating that a critical number of galactose residues was required for efficient uptake by the asialoglycoprotein receptor, similar to the results found for galactosylated BSA and superoxide dismutase (Nishikawa *et al.*, 1995). The more efficient uptake of galactosylated AGP3 (99.5% by 4 h for ≥ 130 galactose moieties) compared to galactose-modified 1E8 was unlikely due to differences in the surface density of galactose groups since galactose incorporation was similar based on the molecular weights of 1E8 and AGP3. The asialoglycoprotein receptor is a hetero-oligomer composed of two types of polypeptide chains (Lodish, 1991). High affinity binding of ligands involves the interaction of three galactose residues in ligands with sites in both polypeptide chains (Rice *et al.*, 1990). Specific orientation of the galactose residues is required for high affinity receptor binding (Lee *et al.*, 1983; Townsend *et al.*, 1986), suggesting that the orientation of incorporated galactose groups in the pentameric structure of IgM may promote high-affinity binding to the asialoglycoprotein receptor. On the basis of the elimination kinetics of the antibodies, 1E8-36G and AGP3-208G were selected for further study.

In vivo Clearance of β G-PEG: The ability of mAbs 1E8 and AGP3 to clear β G-sPEG from the circulation is shown in Figure 9. Mice injected with 200 μ g of β G-sPEG at time 0 were i.v. injected 23 h later with PBS () or 250 μ g of 1E8 (Δ), 1E8-36G (\blacktriangle),

AGP3 (□), or AGP3-208G (■). The concentration of β G-sPEG in serum samples was measured and normalized as described in the Experimental Procedures. Mean concentrations from three mice are shown. Significant differences between AGP3 and 1E8 clearance are indicated by: *, $p \leq 0.05$; **, $p \leq 0.005$; ***, $p \leq 0.0005$. Significant differences between 1E8-36G and AGP3-208G clearance are indicated by: +, $p \leq 0.05$; ++, $p \leq 0.005$; +++, $p \leq 0.0005$. Bars indicate the standard error of the mean. As shown in Figure 9, both 1E8 and AGP3 accelerated the clearance of β G-sPEG from the circulation. 1E8-36G cleared significantly more β G-sPEG than 1E8 at early times but by 8 h after antibody administration there was no significant difference between these antibodies. AGP3 and AGP3-208G cleared significantly more β G-sPEG than either 1E8 or 1E8-36G at all times examined. β G-sPEG was cleared equally well by AGP3 and AGP3-208G under these experimental conditions.

Figure 10 shows a dose dependent relation between AGP3 dose and efficiency of β G-sPEG clearance. Groups of three mice were injected at time 0 with 50 μ g of β G-sPEG followed 24 h later by PBS (●), 150 μ g of AGP3 (□), 150 μ g of AGP3-208G (Δ), 300 μ g of AGP3 (■), or 300 μ g of AGP3-208G (\blacktriangle). The concentration of β G-PEG in serum samples was measured and normalized as described in the Experimental Procedures. Mean concentrations from three mice are shown. Significant differences between AGP3 and AGP3-208G groups are indicated by; *, $p \leq 0.05$. Bars indicate the standard error of the mean. Figure 10 shows that i.v. injection of 150 or 300 μ g of AGP3 or AGP3-208G 24 h after i.v. injection of 50 μ g of β G-sPEG resulted in rapid removal of 90-95% of the enzyme within 1.5 h. The concentration of β G-sPEG in the circulation, however, rebounded from a

minimum of about 2 $\mu\text{g/mL}$ to 4-5 $\mu\text{g/mL}$ 8 h after injection of AGP3-208G. In contrast, the concentration of $\beta\text{G-sPEG}$ decreased to 1.0 $\mu\text{g/mL}$ 8 h after injection of 300 μg of AGP3. Although a dose of 300 μg of AGP3 appeared to more completely clear $\beta\text{G-sPEG}$ compared to 150 μg of AGP3, the difference was only significant ($p \leq 0.05$) 1.5 h after antibody administration. In contrast, clearance of $\beta\text{G-sPEG}$ with both 150 and 300 μg of AGP3 was significantly ($p \leq 0.05$) better than clearance with either dose of AGP3-208G at all times. The better clearance achieved by unmodified AGP3 can be attributed to the prolonged half-life of AGP3 ($\tau_{1/2} = 13$ h) compared to AGP3-208G ($\tau_{1/2} < 1$ h). $\beta\text{G-sPEG}$ released from normal tissues after rapid removal from blood could be bound and cleared by free AGP3 remaining in the blood pool whereas the rapid endocytosis of both free and complexed AGP3-208G prevented further formation of complexes with $\beta\text{G-sPEG}$. This behavior is clearly evident in figure 10 where the concentration of $\beta\text{G-sPEG}$ in the circulation rebounded after an initial period of rapid clearance by AGP3-208G. AGP3 also efficiently cleared 5A8- $\beta\text{G-PEG}$, reducing the serum concentration of conjugate by 280 fold within 2 hours and by 940 fold within 48 hours, as shown in Table 2.

Hindered binding of 1E8 by PEG chains on $\beta\text{G-sPEG}$ may have contributed to the high residual serum concentrations of $\beta\text{G-sPEG}$ after 1E8 administration. The pentameric structure of AGP3 combined with the presence of multiple PEG chains on each βG tetramer may have promoted the formation of large immune complexes, which are cleared more rapidly than small immune complexes. AGP3 appears to produce more rapid and complete clearance than other antibodies previously employed for clearance, which have been limited to IgG fractions of polyclonal antiserum (Bradwell *et al.*, 1983; Sharkey *et al.*, 1984;

Begent *et al.*, 1989; Pedley *et al.*, 1989) or monoclonal IgG antibodies (Sharma *et al.*, 1990; Kerr *et al.*, 1993; Haisma *et al.*, 1995). Polyclonal IgG antibodies reduced radioimmunoconjugate levels in serum from around 10-fold (Bradwell *et al.*, 1983; Sharkey *et al.*, 1984; Begent *et al.*, 1989) to 130-fold (Pedley *et al.*, 1989) in 24 h whereas monoclonal IgG antibodies reduced the serum levels of antibody-enzyme conjugates from 3-10 fold in a few hours (Sharma *et al.*, 1990; Haisma *et al.*, 1995) to 40-60 fold in 24 h (Kerr *et al.*, 1993; Wallace *et al.*, 1994). In comparison, AGP3 reduced the serum concentration of 5A8- β G-PEG by 280-fold in 2 h and 940-fold in 24 h. Although it is difficult to directly compare clearance results employing different immunoconjugates and experimental conditions, these results suggest that IgM may clear immunoconjugates from blood more rapidly and completely than IgG.

Table 2

Clearance of 5A8- β G-PEG by AGP3

Time (hour)	No clearance (μ g/mL)	AGP3 clearance (μ g/mL)	reduction ^b (fold)
24	24 \pm 2.1	37 \pm 1.5	1.0
26	18 \pm 3.5	0.10 \pm 0.001	280
48	6.1 \pm 0.7	0.01 \pm 0	940

^a Groups of 3 mice were i.v. injected with 175 μ g of 5A8- β G-PEG. At 24 and 25 hours, mice were injected with PBS (no clearance) or 300 μ g of AGP3. The concentration of 5A8- β G-PEG in serum was determined by measuring β G activity. ^b The ratio of 5A8- β G-PEG in serum without clearance to the concentration with clearance, normalized for the difference in conjugate concentrations at 24 hour.

PEG modification is a useful technique to improve the stability, prolong serum half-life and decrease the antigenicity and immunogenicity of proteins. Since AGP3 binding to PEG is protein independent, this antibody may be generally applicable to clearance of any PEG-modified immunoconjugates. However, clearance of immunoconjugates from the circulation may also result in decreased tumor accumulation of the immunoconjugates, probably due to a concentration gradient of the conjugates from tumor to blood, which allows the immunoconjugates to be dissociated from tumor antigen and released into the blood. But this undesirable process may be slowed down by using IgM type antibodies since IgM is larger in size, which may delay its entrance into the tumor interstitial space thereby minimizing interactions between localized immunoconjugates and the antibody. In addition, the rapid clearance achieved by using AGP3 permits administration of prodrugs or other targets before a significant amount of immunoconjugates are released from the tumor.

Prodrug Toxicity after Clearance of β G-sPEG: Clearance of circulating β G-sPEG should reduce systemic activation of the glucuronide prodrug BHAMG, thereby reducing toxicity. One concern of employing IgM to clear proteins is that the immune complexes could be deposited in organs and cause tissue damage. In addition, deposition of β G-PEG on the surface of cells could result in prodrug activation and tissue toxicity. Organ tissue sections from mice that were injected with PBS or β G-sPEG followed 24 h later by AGP3 and then BHAMG were examined after 3 or 10 days as shown in Figure 11. Mice were injected with PBS (A-C) or β G-sPEG (D-F) on day 1. Mice were i.v. injected with AGP3 and with BHAMG 24 and 30 h later, respectively. Tissue sections of liver (A, D), kidney

(B, E), and spleen (C, F) recovered 10 days later were stained with hematoxylin and eosin.

Figure 11 shows that the injection of AGP3 and BHAMG as well as clearance of β G-sPEG with AGP3 before BHAMG administration did not cause obvious damage to the liver, spleen, or kidney 10 days after BHAMG administration. Similarly, no tissue damage was observed 3 days after prodrug administration (results not shown).

The toxicity associated with clearance and prodrug administration was also assessed by measuring the weight and blood cells of mice, as shown in Figures 12A and 12B. Groups of 4 mice were i.v. injected at time zero with PBS (control) or β G-sPEG. Twenty-four hours later, mice received i.v. injections of 300 μ g of AGP3 (control and AGP3 groups), 300 μ g of AGP3-208G, 150 μ g of AGP3-208G (x2, 4 h delay), or PBS (no clearance). Six hours later, all mice were i.v. injected with BHAMG. Figure 12A shows the time course of mouse weight (% of initial weight) and Figure 12B shows the mean numbers of WBC, RBC, and platelets on day 8. Significant differences between the control and experimental groups are indicated by: *, $p \leq 0.05$; **, $p \leq 0.005$; ***, $p \leq 0.0005$. Bars indicate the standard error of the mean. Administration of AGP3 and 15 mg/kg BHAMG (control group) produced only a small transient decrease in the weight of mice (Figure 12A). In contrast, injection of 50 μ g of β G-sPEG 24 h before administration of BHAMG (no clearance) caused significant weight loss and resulted in the death of 1 of 4 mice. Clearance of β G-sPEG with 300 μ g of AGP3 or 2 injection of 150 μ g of AGP3-208G before prodrug administration prevented significant weight loss compared to the control group. Mice that were cleared with one 300 μ g dose of AGP3-208G before injection of BHAMG lost significantly more weight than control mice after day 8, but the

loss was much less than in the uncleared mice. Figure 12B shows that the numbers of RBC, WBC and PLT were significantly reduced in mice injected with β G-sPEG and BHAMG (no clearance) compared to mice injected with AGP3 and BHAMG (control mice). In contrast, the numbers of RBC, WBC, and PLT in mice in which β G-sPEG was
5 cleared with 300 μ g of AGP3 before BHAMG administration were not significantly different from control mice. Clearance of β G-sPEG with AGP3-208G before injection of prodrug reduced but did not eliminate hematological toxicity.

AGP3/ β G-PEG immune complexes may be routed to the acidic environment of lysosomes after clearance. The pH sensitivity of β G is shown in Figure 13. β G was assayed
10 for enzymatic activity at various pH values (\square) or incubated at the indicated pH for 4 h before neutralization and assay at pH 7.0 (O). Results represent percentage of maximum activity. β G was irreversibly deactivated at pH values less than 5.0 or greater than 9.0 (Figure 13).

Soluble IgM immune complexes are primarily removed from the circulation and
15 catabolized by the mononuclear phagocyte system in the liver, spleen and lungs, possibly by receptor-mediated binding of high mannose oligosaccharides exposed upon conformational changes in IgM after binding to antigen (Day *et al.*, 1980). The low toxicity of BHAMG after clearance of β G-sPEG with AGP3 indicates that immune complexes formed between AGP3 and β G-sPEG are rapidly internalized where β G is
20 inactivated, degraded or inaccessible to prodrug. Routing of β G-sPEG to lysosomes is expected to result in loss of enzymatic activity because β G is irreversibly inactivated at pH values lower than 5.0. Deactivation of β G in lysosomes may prevent accumulation of

active enzyme and prodrug activation in the liver as observed after antibody-mediated clearance of carboxypeptidase G2 conjugates (Rogers *et al.*, 1995).

Characterization of the LS174T tumor model: LS174T colon adenocarcinoma cells were used as a human tumor model to examine the effect of clearance on the localization and therapeutic efficacy of glucuronide prodrug activation by B72.3- β G-PEG, a conjugate formed by covalently linking F(ab')₂ fragments of mAb B72.3 to PEG-modified β G. In Figure 14A, mAb B72.3 (\square), B72.3- β G-PEG (Δ) and the control conjugate H25- β G-PEG (O) were assayed by ELISA for binding to bovine submaxillary gland mucin. Absorbance (405 nm) of wells was measured 30 min after addition of ABTS substrate. In Figure 14B, the absorbance (405 nm) of wells containing the indicated concentrations of β G (\square), B72.3- β G-PEG (Δ) and H25- β G-PEG (O) was measured 15 min after addition of PNPG substrate. Results show the mean values of duplicate determinations. Bars indicate the standard error of the mean. B72.3- β G-PEG retained 75% of the mucin binding activity of mAb B72.3 (Fig. 14A) and 100% of the enzyme activity of unmodified β G (Fig. 14B). Control conjugate H25- β G-PEG did not bind LS174T cells (Fig. 14A) but retained 92.4% of native β G activity (Fig. 14B).

As shown in Figure 15, frozen section of LS174T tumors were incubated with mAb B72.3 (A) or H25B10 (B) followed by biotin-labeled goat anti-mouse Ig and streptavidin-HRP before addition of substrate. Magnification is 200x. The sections were counterstained with hematoxylin. Examples of positive immunostaining are indicated with arrows. Immunohistochemical analysis of antibody binding confirmed that mAb B72.3 (Fig. 15A) but not mAb H25B10 (Fig. 15B) bound to LS174T xenografts. mAb B72.3 binding to

LS174T sections was heterogeneous (Fig. 15A). The sensitivity of LS174T tumor cells to pHAM and BHAMG was determined by measuring [^3H]-leucine incorporation into cellular proteins after exposure to drugs for 24 h. Comparison of IC_{50} values showed that BHAMG was 800 times less toxic than pHAM to LS174T cells as shown in Figure 16 where

5 LS174T colon adenocarcinoma cells were exposed to pHAM (\square), BHAMG (O) or BHAMG plus 1 $\mu\text{g}/\text{well}$ βG (Δ) for 24 h, washed with PBS and incubated in fresh medium for 24 h before [^3H]-leucine incorporation was determined. The cellular protein synthesis rate of drug-treated cells is compared to untreated control cells. Bars indicate the standard error of triplicate determinations. The simultaneous addition of βG and BHAMG to tumor

10 cells resulted in a cytotoxic effect equal to pHAM alone, indicating efficient cleavage of the glucuronide functional group of BHAMG.

Tumor localization of B72.3- βG -PEG: Figure 17 shows that ^{125}I -B72.3- βG -PEG preferentially localized in LS174T xenografts. Radioactivity of tumors and tissues was determined (A) 24, (B) 48, (C) 72 and (D) 96 h after nude mice bearing 100-200 mm^3

15 LS174T tumors were i.v. injected with 200 μg (140 μCi) ^{125}I -B72.3- βG -PEG (solid columns) or ^{125}I -H25- βG -PEG (open columns). Results represent the mean values of three mice. Significant differences between B72.3- βG -PEG and H25- βG -PEG are indicated by; *, $p \leq 0.05$; **, $p \leq 0.005$. Bars show the standard error of the mean. Maximum accumulation of ^{125}I -B72.3- βG -PEG in LS174T xenografts ($3.11 \pm 0.8\%$ injected dose/g)

20 was achieved within 48 h after injection. However, the tumor to blood ratio at 48 h was only 1.9. The tumor/blood ratio of B72.3- βG -PEG increased to 4.8 at 96 h but

immunoenzyme in tumors decreased about 3-fold from levels achieved at 48 h. ^{125}I -H25- β G-PEG did not specifically localize in LS174T xenografts (Figs. 17A-D).

In vivo clearance of B72.3- β G-PEG and H25- β G-PEG: The ability of AGP3 to clear B72.3- β G-PEG and H25- β G-PEG from the circulation was examined by i.v. injecting BALB/c mice with 250 μg conjugates at time 0 followed 48 and 50 h later by two i.v. injections of 300 and 200 μg AGP3 or PBS. Mean serum concentrations from 2-3 mice are shown. Bars indicate the standard error of the mean. Figure 18 shows that AGP3 reduced the concentration of B72.3- β G-PEG in blood by 33-fold (13.4 to 0.41 $\mu\text{g}/\text{ml}$) and H25- β G-PEG by 65-fold (16.8 to 0.26 $\mu\text{g}/\text{ml}$) in 6 h.

Tumor localization of B72.3- β G-PEG with AGP3 clearance: The effect of clearance on tumor localization was determined in groups of 6-7 BALB/c *nu/nu* mice bearing 100-200 mm^3 LS174T tumor xenografts. Mice were i.v. injected at time 0 with 200 μg (140 μCi) ^{125}I -B72.3- β G-PEG (Fig. 19A) or ^{125}I -H25- β G-PEG (Fig. 19B). At 48 and 50 h, half the mice were i.v. injected with two fractionated doses of AGP3 (300 and 200 μg , open columns) whereas the other half received PBS (solid columns). Mice were sacrificed after 6 h and tumors, blood and organs were assayed for radioactivity. Results represent the mean values of 3-4 mice. Significant differences between immunoenzyme uptake with and without AGP3 clearance are indicated by; *, $p \leq 0.05$; **, $p \leq 0.005$. Bars show the standard error of the mean. Figure 19A shows that tumor localization of B72.3- β G-PEG after AGP3 clearance did not significantly ($p > 0.1$) differ from tumor uptake without clearance (2.35 ± 0.18 versus 2.14 ± 0.23 % injected dose / g). In contrast, the radioactivity in serum and all tissues except for the colon was lower with AGP3 clearance. This resulted in improved

tumor/tissue ratios of B72.3- β G-PEG for most tissues (Table 3). For example, the tumor/blood ratio increased from 3.9 ± 0.2 to 29.6 ± 3.1 . Although the control conjugate did not specifically accumulate in LS174T xenografts, clearance with AGP3 further decreased the concentration of H25- β G-PEG in the blood and most tissues (Figure 19B).

5 Thus, AGP3 not only reduced the serum concentration of conjugates by 33-65-fold in 6 h but also accelerated the clearance of immunoenzyme from most normal tissues. Importantly, AGP3 did not significantly reduce tumor accumulation of B72.3- β G-PEG, in contrast to some other clearance systems (Pedley *et al.*, 1989; Sharkey *et al.*, 1992; Kerr *et al.*, 1993). The large size of AGP3 (IgM) may hinder passage of the mAb into the tumor interstitial
10 space, minimizing interactions between localized immunoconjugate and AGP3. The low tumor/blood ratio without clearance can be attributed to the loss of immunoenzyme from the tumor during the prolonged period required for immunoenzyme to reach safe levels in serum. Clearance of immunoenzyme with AGP3 can allow earlier administration of prodrug when immunoenzyme localization at tumor cells is maximal.

Table 3

The tumor/tissues ratio of conjugates after clearance with AGP3

Tissue	Tumor / Tissue Ratio			
	B72.3- β G-PEG		H25- β G-PEG	
	No clearance	AGP3 clearance	No clearance	AGP3 clearance
Tumor	1	1	1	1
Blood	3.9 ± 0.2	29.6 ± 3.1	0.3 ± 0.1	2.2 ± 0.8
Lung	12.5 ± 3.3	33.5 ± 3.0	1.4 ± 0.4	5.5 ± 2.1
Liver	15.9 ± 2.9	19.2 ± 0.8	1.1 ± 0.2	1.5 ± 0.6
Spleen	19.0 ± 2.6	27.7 ± 5.2	1.8 ± 0.5	3.5 ± 1.5
Kidney	18.1 ± 1.3	31.0 ± 2.0	1.4 ± 0.5	2.2 ± 0.2
Intestine	6.4 ± 1.7	165 ± 56	4.8 ± 1.3	10.1 ± 2.9
Colon	54.3 ± 7.4	50.2 ± 8.3	3.6 ± 0.5	3.1 ± 0.33
Urine	1.1 ± 0.1	0.6 ± 0.1	0.1 ± 0.02	0.03 ± 0.001

Groups of 3 or 4 mice bearing 100-200 mm³ solid LS174T tumors were injected with radiolabeled conjugates follow 48 h later by two i.v. injections of AGP3 or PBS. After 6 h, tumors, blood, urine and normal tissues were removed and counted in a gamma counter.

The radioactivity in urine after clearance of both B72.3- β G-PEG and H25- β G-PEG with AGP3 was significantly ($p \leq 0.005$) greater than without clearance (results not shown).

Autoradiographic analysis of urine samples after electrophoresis on SDS PAGE revealed the

presence of low-molecular-weight ^{125}I -labeled peptides but no intact conjugates (data not shown), indicating that AGP3 clearance resulted in the rapid metabolism of conjugates.

Therapy of LS174T xenografts: The antitumor activity of BHAMG in combination with B72.3- βG -PEG after clearance of free conjugate with AGP3 was examined. As shown in Figure 20A, groups of 9 BALB/c *nu/nu* mice bearing 50-100 mm³ LS174T tumors were i.v. injected with B72.3- βG -PEG (□) or H25- βG -PEG (■) on day 9 followed by two i.v. injections of AGP3 on day 11. After 6 h, mice were i.v. injected with BHAMG (7.5 mg/kg x 3). Control groups of tumor-bearing mice were treated with BHAMG (Δ), pHAM (O) or PBS (●) alone. Therapy was repeated starting on days 16, 26 and 41. Results show the mean tumor size. Bars indicate the standard error of the mean. The mean size of tumors in mice sequentially treated with B72.3- βG -PEG, AGP3 and BHAMG was significantly ($p \leq 0.0005$) smaller than control tumors after day 14. Table 4 shows that treatment toxicity with clearance was minimal with a maximum weight loss of 6 % over 4 rounds of therapy. In contrast, pHAM treatment caused a maximum weight loss of 13% over 2 rounds of therapy even though it did not provide antitumor activity. As shown in Figure 20B, groups of 8 nude mice bearing larger (200-250 mm³) LS174T tumors were treated with two rounds of therapy as above starting on days 11 and 23. The mean size of tumors in mice sequentially treated with B72.3- βG -PEG, AGP3 and BHAMG was significantly ($p \leq 0.005$) smaller than control tumors after day 16. The degree of tumor suppression achieved with minimal toxicity by prodrug therapy compares favorably with conventional drugs such as 5-fluorouracil (Blumenthal *et al.*, 1994) and doxorubicin (Meyer *et al.*, 1995) in the LS174T xenograft model. B72.3- βG -PEG appeared to be rapidly catabolized into

small peptides that were eliminated in the urine and possibly bile. Soluble IgM immune complexes are primarily removed from the circulation and catabolized by the mononuclear phagocyte system in the liver, spleen and lungs by receptor-mediated binding of high mannose oligosaccharides exposed upon conformational changes in IgM induced by antigen binding (Day *et al.*, 1980). The low toxicity of BHAMG treatment observed after clearance of B72.3- β G-PEG with AGP3 indicates that cleared conjugate was unavailable for prodrug activation, consistent with rapid degradation of the immunoenzyme after clearance.

Table 4

Toxicity of therapy

	Weight loss %			
	Round of therapy			
	1	2	3	4
PBS	0	0	0	NM
pHAM	6.8	13.0	NM	NM
BHAMG	1.7	4.0	1.7	-
B72.3- β G-PEG / AGP3 / BHAMG	3.5	5.0	3.8	6.0
H25- β G-PEG / AGP3 / BHAMG	3.9	6.6	4.6	NM

NM: not meaningful due to mouse deaths.

5

10

Most of the enzymes currently under investigation for the targeted activation of anti-neoplastic prodrugs are of microbial origin such as β -lactamase from *Enterobacter cloacae* (Kerr *et al.*, 1995), carboxypeptidase G2 from *Pseudomonas* species (Blakey *et al.*, 1996), β -glucuronidase (Chen *et al.*, 1997), nitroreductase (Anlezark *et al.*, 1995) and penicillin-G amidase (Vrudhula *et al.*, 1993) from *Escherichia coli*, and cytosine deaminase from bakers yeast (Wallace *et al.*, 1994). These enzymes are expected to induce a strong immune response as has been found for carboxypeptidase G2 in a pilot clinical trial (Sharma *et al.*, 1996), thereby limiting the number of times that immunoenzymes can be administered to

patients. The utilization of human enzymes to activate antineoplastic prodrugs may not completely prevent this problem since recombinant human proteins can also induce immune responses in patients (Atkins *et al.*, 1986; Gribben *et al.*, 1990). Immunosuppressive drugs such as cyclosporin A, cyclophosphamide and deoxyspergualin can decrease or delay the immune response against antibodies (Ledermann *et al.*, 1988), immunotoxins (Pai *et al.*, 1990) and antibody-enzyme conjugates (Sharma *et al.*, 1996). Immunosuppression, in addition to producing toxicity in some patients (Sharma *et al.*, 1996), may hinder the development of antitumor immunity generated by prodrug therapy (Chen *et al.*, 1997).

Immune responses against proteins can be attenuated by attachment of PEG (Abuchowski *et al.*, 1977). PEG, in contrast to immunosuppressive drugs, is not toxic and does not affect systemic immunity. PEG modification has been shown to reduce the immunogenicity of enzymes (Abuchowski *et al.*, 1977b), antibodies (Kitamura *et al.*, 1991), toxins (Wang *et al.*, 1993) and recombinant human proteins (Katre, 1990). PEG-modification of bacterial enzymes may allow repeated administration of immunoenzymes for ADEPT without the need to employ toxic immunosuppressive drugs. PEG-modified enzymes of the proper molecular size can also preferentially accumulate in tumors and are under investigation for tumor selective prodrug activation (Bagshawe *et al.*, 1999). AGP3 binds to the backbone of PEG independent of the linker or protein employed. AGP3 should therefore be generally applicable to the accelerated clearance of PEG-modified immunoenzymes, radioimmunoconjugates and imaging agents as well as for the analysis of PEG-modified proteins.

The following references are cited to further explain or describe the present invention,
the contents of which are incorporated by reference.

- 5 (1) Abuchowski A, McCoy JR, Palczuk NC, van Es T and Davis FF (1977a) Effect of
covalent attachment of polyethylene glycol on immunogenicity and circulating life of
bovine liver catalase. J-Biol-Chem 252, 3582-3586.
- (2) Abuchowski A, van Es T, Palczuk NC and Davis FF (1977b) Alteration of immunological
properties of bovine serum albumin by covalent attachment of polyethylene glycol. J-
Biol-Chem 252, 3578-3581.
- 10 (3) Anlezark GM, Melton RG, Sherwood RF, Wilson WR, Denny WA, Palmer BD, Knox RJ,
Friedlos F and Williams A (1995) Bioactivation of dinitrobenzamide mustards by an E.
coli B nitroreductase. Biochem Pharmacol 50, 609-618.
- (4) Atkins MB, Gould JA, Allegretta M, Li JJ, Dempsey RA, Rudders RA, Parkinson DR,
Reichlin S and Mier JW (1986) Phase I evaluation of recombinant interleukin-2 in
15 patients with advanced malignant disease. J-Clin-Oncol 4, 1380-1391.
- (5) Bagshawe KD, Sharma SK, Know R and Amooquaye E (1999) Studies with ploymer
enzyme conjugates and AMIRACS. Advances in the Application of Monoclonal
Antibodies in Clinical Oncology Samos, Greece.

(6) Bagshawe KD, Springer CJ, Searle F, Antoniow P, Sharma SK, Melton RG and Sherwood RF (1988) A cytotoxic agent can be generated selectively at cancer sites. Br J Cancer 58, 700-703.

(7) Bassar RL, Rasko JE, Clarke K, Cebon J, Green MD, Hussein S, Alt C, Menchaca D, Tomita D, Marty J, Fox RM and Begley CG (1996) Thrombopoietic effects of pegylated recombinant human megakaryocyte growth and development factor (PEG-rHuMGDF) in patients with advanced cancer. Lancet 348, 1279-1281.

(8) Beatty JD, Beatty BG and Vlahos WG (1987) Measurement of monoclonal antibody affinity by non-competitive enzyme immunoassay. J Immunol Methods 100, 173-179.

(9) Beckman JS, Minor RL, Jr., White CW, Repine JE, Rosen GM and Freeman BA (1988) Superoxide dismutase and catalase conjugated to polyethylene glycol increases endothelial enzyme activity and oxidant resistance. J-Biol-Chem 263, 6884-6892.

(10) Begent RH, Ledermann JA, Green AJ, Bagshawe KD, Riggs SJ, Searle F, Keep PA, Adam T, Dale RG and Glaser MG (1989) Antibody distribution and dosimetry in patients receiving radiolabelled antibody therapy for colorectal cancer. Br J Cancer 60, 406-412.

(11) Blakey DC, Burke PJ, Davies DH, Dowell RI, East SJ, Eckersley KP, Fitton JE, McDaid J, Melton RG, Niculescu Duvaz IA, Pinder PE, Sharma SK, Wright AF and Springer CJ (1996) ZD2767, an improved system for antibody-directed enzyme prodrug therapy that results in tumor regressions in colorectal tumor xenografts. Cancer Res 56, 3287-3292.

- (12) Blumenthal RD, Sharkey RM, Natale AM, Kashi R, Wong G and Goldenberg DM (1994)

Comparison of equitoxic radioimmunotherapy and chemotherapy in the treatment of human colonic cancer xenografts. Cancer-Res 54, 142-151.

- (13) Bosslet K, Czech J, Seemann G, Monneret C and Hoffmann D (1994) Fusion protein

mediated prodrug activation (FMPA) in vivo. Cell Biophys 24-25, 51-63.

- (14) Bradwell AR, Vaughan A, Fairweather DS and Dykes PW (1983) Improved

radioimmunodetection of tumours using second antibody [letter]. Lancet 1, 247.

- (15) Brinckerhoff LH, Kalashnikov VV, Thompson LW, Yamshchikov GV, Pierce RA,

Galavotti HS, Engelhard VH and Slingluff CL, Jr. (1999) Terminal modifications inhibit

proteolytic degradation of an immunogenic MART-1(27-35) peptide: implications for peptide vaccines. Int-J-Cancer 83, 326-334.

- (16) Cabanes A, Tzemach D, Goren D, Horowitz AT and Gabizon A (1998) Comparative

study of the antitumor activity of free doxorubicin and polyethylene glycol-coated liposomal doxorubicin in a mouse lymphoma model. Clin-Cancer-Res 4, 499-505.

- (17) Caliceti P, Monfardini C, Sartore L, Schiavon O, Baccichetti F, Carllassare F and

Veronese FM (1993) Preparation and properties of monomethoxy poly(ethylene glycol) doxorubicin conjugates linked by an amino acid or a peptide as spacer. Farmaco 48, 919-932.

- (18) Chaffee S, Mary A, Stiehm ER, Girault D, Fischer A and Hershfield MS (1992) IgG antibody response to polyethylene glycol-modified adenosine deaminase in patients with adenosine deaminase deficiency. J-Clin-Invest 89, 1643-1651.
- (19) Chapman AP, Antoniwi P, Spitali M, West S, Stephens S and King DJ (1999) Therapeutic antibody fragments with prolonged in vivo half-lives. Nat-Biotechnol 17, 780-783.
- (20) Chen BM, Chan LY, Wang SM, Wu MF, Chern JW and Roffler SR (1997) Cure of malignant ascites and generation of protective immunity by monoclonal antibody-targeted activation of a glucuronide prodrug in rats. Int J Cancer 73, 392-402.
- (21) Chen RH, Abuchowski A, Van Es T, Palczuk NC and Davis FF (1981) Properties of two urate oxidases modified by the covalent attachment of poly(ethylene glycol). Biochim-Biophys-Acta 660, 293-298.
- (22) Cheng TL, Chen BM, Chan LY, Wu PY, Chern JW and Roffler SR (1997) Poly(ethylene glycol) modification of beta-glucuronidase-antibody conjugates for solid-tumor therapy by targeted activation of glucuronide prodrugs. Cancer-Immunol-Immunother 44, 305-315.
- (23) Cheng TL, Chen BM, Chern JW, Wu MF and Roffler SR (In press) Efficient clearance of poly(ethylene glycol)-modified immunoenzyme with anti-PEG monoclonal antibody for prodrug cancer therapy. Bioconj Chem, .

(24) Cheng TL, Wu PY, Wu MF, Chern JW and Roffler SR (1999) Accelerated clearance of polyethylene glycol-modified proteins by anti-polyethylene glycol IgM. Bioconjug-Chem 10, 520-528.

5 (25) Childs CE (1975) The determination of polyethylene glycol in gamma globulin solutions. Microchem-J 20, 190-192.

(26) Chinol M, Casalini P, Maggiolo M, Canevari S, Omodeo ES, Caliceti P, Veronese FM, Cremonesi M, Chiolerio F, Nardone E, Siccardi AG and Paganelli G (1998) Biochemical modifications of avidin improve pharmacokinetics and biodistribution, and reduce immunogenicity. Br-J-Cancer 78, 189-197.

10 (27) Conover CD, Pendri A, Lee C, Gilbert CW, Shum KL and Greenwald RB (1997) Camptothecin delivery systems: the antitumor activity of a camptothecin-20-0-polyethylene glycol ester transport form. Anticancer-Res 17, 3361-3368.

15 (28) Day JF, Thornburg RW, Thorpe SR and Baynes JW (1980) Carbohydrate-mediated clearance of antibody . antigen complexes from the circulation. The role of high mannose oligosaccharides in the hepatic uptake of IgM . antigen complexes. J Biol Chem 255, 2360-2365.

(29) Delgado C, Francis GE and Fisher D (1992) The uses and properties of PEG-linked proteins. Crit Rev Ther Drug Carrier Syst 9, 249-304.

20 (30) Delgado C, Pedley RB, Herraiz A, Boden R, Boden JA, Keep PA, Chester KA, Fisher D, Begent RH and Francis GE (1996) Enhanced tumour specificity of an anti-

carcinoembrionic antigen Fab' fragment by poly(ethylene glycol) (PEG) modification. Br J Cancer 73, 175-182.

(31) Divgi CR, Scott AM, Fallone PS, Hilton S, Siler K, Finn RD and Cohen AM (1994) Clinical comparison of radiolocation of two monoclonal antibodies against the TAG-72 antigen. Nucl Med Biol 21, 9-15.

(32) Eccles SA, Court WJ, Box GA, Dean CJ, Melton RG and Springer CJ (1994) Regression of established breast carcinoma xenografts with antibody-directed enzyme prodrug therapy against c-erbB2 p185. Cancer Res 54, 5171-5177.

(33) Esslinger HU, Haas S, Maurer R, Lassmann A, Dubbers K and Muller Peltzer H (1997) Pharmacodynamic and safety results of PEG-Hirudin in healthy volunteers. Thromb-Haemost 77, 911-919.

(34) Fareed J, Callas D, Hoppensteadt DA, Walenga JM and Bick RL (1998) Antithrombin agents as anticoagulants and antithrombotics. Implications in drug development. Med-Clin-North-Am 82, 569-586.

(35) Goffin V, Bernichtein S, Carriere O, Bennett WF, Kopchick JJ and Kelly PA (1999) The human growth hormone antagonist B2036 does not interact with the prolactin receptor. Endocrinology 140, 3853-3856.

(36) Greenwald RB, Pendri A, Conover CD, Lee C, Choe YH, Gilbert C, Martinez A, Xia J, Wu D and Hsue M (1998) Camptothecin-20-PEG ester transport forms: the effect of spacer groups on antitumor activity. Bioorg-Med-Chem 6, 551-562.

(37) Gribben JG, Devereux S, Thomas NS, Keim M, Jones HM, Goldstone AH and Linch DC

(1990) Development of antibodies to unprotected glycosylation sites on recombinant human GM-CSF [see comments]. Lancet 335, 434-437.

(38) Guermant C, Brygier J, Baeyens Volant D, Nijs M, Vincentelli J, Paul C and Looze Y

5 (1995) Quantitative determination of polyethylene glycol based upon its salting out and partitioning of a dye into the resulting aqueous two-phase system. Anal-Biochem 230, 254-258.

(39) Habeeb AF (1966) Determination of free amino groups in proteins by trinitrobenzenesulfonic acid. Anal Biochem 14, 328-336.

10 (40) Haisma HJ, Van Muijen M, Scheffer G, Scheper RJ, Pinedo HM and Boven E (1995) A monoclonal antibody against human beta-glucuronidase for application in antibody-directed enzyme prodrug therapy. Hybridoma 14, 377-382.

(41) Harker LA (1999) Physiology and clinical applications of platelet growth factors. Curr-Opin-Hematol 6, 127-134.

15 (42) He XH, Shaw PC and Tam SC (1999) Reducing the immunogenicity and improving the in vivo activity of trichosanthin by site-directed pegylation. Life Sci 65, 355-368.

(43) Hudson PJ (1998) Recombinant antibody fragments. Curr-Opin-Biotechnol 9, 395-402.

(44) Ito HO, So T, Hirata M, Koga T, Ueda T and Imoto T (1998) Tolerogenic activity of polyethylene glycol-conjugated lysozyme distinct from that of the native counterpart.

20 Immunology 93, 200-207.

- (45) Ito HO, So T, Ueda T, Imoto T and Koga T (1997) Prevention of collagen-induced arthritis (CIA) by treatment with polyethylene glycol-conjugated type II collagen; distinct tolerogenic property of the conjugated collagen from the native one. Clin-Exp-Immunol 108, 213-219.
- 5 (46) Johnson VG, Schlom J, Paterson AJ, Bennett J, Magnani JL and Colcher D (1986) Analysis of a human tumor-associated glycoprotein (TAG-72) identified by monoclonal antibody B72.3. Cancer Res 46, 850-857.
- (47) Kaneda Y, Yamamoto S, Kihira T, Tsutsumi Y, Nakagawa S, Miyake M, Kawasaki K and Mayumi T (1995) Synthetic cell-adhesive laminin peptide YIGSR conjugated with
10 polyethylene glycol has improved antimetastatic activity due to a longer half-life in blood. Invasion Metastasis 15, 156-162.
- (48) Katre NV (1990) Immunogenicity of recombinant IL-2 modified by covalent attachment of polyethylene glycol. J-Immunol 144, 209-213.
- (49) Katre NV, Knauf MJ and Laird WJ (1987) Chemical modification of recombinant
15 interleukin 2 by polyethylene glycol increases its potency in the murine Meth A sarcoma model. Proc-Natl-Acad-Sci-U-S-A 84, 1487-1491.
- (50) Kerr DE, Garrigues US, Wallace PM, Hellstrom KE, Hellstrom I and Senter PD (1993) Application of monoclonal antibodies against cytosine deaminase for the in vivo clearance of a cytosine deaminase immunoconjugate. Bioconjug Chem 4, 353-357.

- (51) Kerr DE, Schreiber GJ, Vrudhula VM, Svensson HP, Hellstrom I, Hellstrom KE and Senter PD (1995) Regressions and cures of melanoma xenografts following treatment with monoclonal antibody beta-lactamase conjugates in combination with anticancer prodrugs. Cancer Res 55, 3558-3563.
- 5 (52) Kinahan IM and Smyth MR (1991) High-performance liquid chromatographic determination of PEG 600 in human urine. J-Chromatogr 565, 297-307.
- (53) King DJ, Turner A, Farnsworth AP, Adair JR, Owens RJ, Pedley RB, Baldock D, Proudfoot KA, Lawson AD, Beeley NR and et al. (1994) Improved tumor targeting with chemically cross-linked recombinant antibody fragments. Cancer Res 54, 6176-6185.
- 10 (54) Kirpotin D, Park JW, Hong K, Zalipsky S, Li WL, Carter P, Benz CC and Papahadjopoulos D (1997) Sterically stabilized anti-HER2 immunoliposomes: design and targeting to human breast cancer cells in vitro. Biochemistry 36, 66-75.
- (55) Kitamura K, Takahashi T, Yamaguchi T, Noguchi A, Noguchi A, Takashina K, Tsurumi H, Inagake M, Toyokuni T and Hakomori S (1991) Chemical engineering of the
15 monoclonal antibody A7 by polyethylene glycol for targeting cancer chemotherapy. Cancer Res 51, 4310-4315.
- (56) Klibanov AL, Martynov AV, Slinkin MA, Sakharov I, Smirnov MD, Muzykantov VR, Danilov SM and Torchilin VP (1988) Blood clearance of radiolabeled antibody: enhancement by lactosamination and treatment with biotin-avidin or anti-mouse IgG
20 antibodies. J Nucl Med 29, 1951-1956.

(57) Kobayashi H, Sakahara H, Endo K, Yao ZS, Toyama S and Konishi J (1995) Repeating the avidin "chase" markedly improved the biodistribution of radiolabelled biotinylated antibodies and promoted the excretion of additional background radioactivity. Eur J Cancer 31a, 1689-1696.

5 (58) Ledermann JA, Begent RH, Bagshawe KD, Riggs SJ, Searle F, Glaser MG, Green AJ and Dale RG (1988) Repeated antitumour antibody therapy in man with suppression of the host response by cyclosporin A. Br J Cancer 58, 654-657.

10 (59) Lee WY, Sehon AH and Akerblom E (1981) Suppression of reaginic antibodies with modified allergens. IV. Induction of suppressor T cells by conjugates of polyethylene glycol (PEG) and monomethoxy PEG with ovalbumin. Int Arch Allergy Appl Immunol 64, 100-114.

(60) Lee YC, Townsend RR, Hardy MR, Lonngren J, Arnarp J, Haraldsson M and Lonn H (1983) Binding of synthetic oligosaccharides to the hepatic Gal/GalNAc lectin. Dependence on fine structural features. J Biol Chem 258, 199-202.

15 (61) Lin L (1998) Betaseron. Dev Biol Stand 96, 97-104.

(62) Lodish HF (1991) Recognition of complex oligosaccharides by the multi-subunit asialoglycoprotein receptor. Trends Biochem Sci 16, 374-377.

20 (63) Maiti PK, Lang GM and Sehon AH (1988) Tolerogenic conjugates of xenogeneic monoclonal antibodies with monomethoxypolyethylene glycol. I. Induction of long-lasting tolerance to xenogeneic monoclonal antibodies. Int J Cancer Suppl 3, 17-22.

- (64) Maloney DG, Kaminski MS, Burowski D, Haimovich J and Levy R (1985) Monoclonal anti-idiotypic antibodies against the murine B cell lymphoma 38C13: characterization and use as probes for the biology of the tumor in vivo and in vitro. Hybridoma 4, 191-209.
- (65) Marshall D, Pedley RB, Melton RG, Boden JA, Boden R and Begent RH (1995) Galactosylated streptavidin for improved clearance of biotinylated intact and F(ab')₂ fragments of an anti-tumour antibody. Br J Cancer 71, 18-24.
- (66) Martin J, Stribbling SM, Poon GK, Begent RH, Napier M, Sharma SK and Springer CJ (1997) Antibody-directed enzyme prodrug therapy: pharmacokinetics and plasma levels of prodrug and drug in a phase I clinical trial. Cancer Chemother Pharmacol 40, 189-201.
- (67) Menzel T, Schomburg A, Korfer A, Hadam M, Meffert M, Dallmann I, Casper S, Kirchner H, Poliwoda H and Atzpodien J (1993) Clinical and preclinical evaluation of recombinant PEG-IL-2 in human. Cancer-Biother 8, 199-212.
- (68) Meyer DL, Law KL, Payne JK, Mikolajczyk SD, Zarrinmayeh H, Jungheim LN, Kling JK, Shepherd TA and Starling JJ (1995) Site-specific prodrug activation by antibody-beta-lactamase conjugates: preclinical investigation of the efficacy and toxicity of doxorubicin delivered by antibody directed catalysis. Bioconjug Chem 6, 440-446.
- (69) Meyer O, Kirpotin D, Hong K, Sternberg B, Park JW, Woodle MC and Papahadjopoulos D (1998) Cationic liposomes coated with polyethylene glycol as carriers for oligonucleotides. J-Biol-Chem 273, 15621-15627.

- (70) Milenic DE, Esteban JM and Colcher D (1989) Comparison of methods for the generation of immunoreactive fragments of a monoclonal antibody (B72.3) reactive with human carcinomas. J Immunol Methods 120, 71-83.
- 5 (71) Miles PJ, Langley KV, Stacey CJ and Talarico TL (1997) Detection of residual polyethylene glycol derivatives in pyridoxylated-hemoglobin-polyoxyethylene conjugate. Artif-Cells-Blood-Substit-Immobil-Biotechnol 25, 315-326.
- (72) Mullin JM, Marano CW, Laughlin KV, Nuciglio M, Stevenson BR and Soler P (1997) Different size limitations for increased transepithelial paracellular solute flux across phorbol ester and tumor necrosis factor-treated epithelial cell sheets. J-Cell-Physiol 171,
10 226-233.
- (73) Nag A, Mitra G and Ghosh PC (1996) A colorimetric assay for estimation of polyethylene glycol and polyethylene glycolated protein using ammonium ferrothiocyanate. Anal-Biochem 237, 224-231.
- (74) Nag A, Mitra G and Ghosh PC (1997) A colorimetric estimation of polyethyleneglycol-conjugated phospholipid in stealth liposomes. Anal-Biochem 250, 35-43.
15
- (75) Nemunaitis J (1997) A comparative review of colony-stimulating factors. Drugs 54, 709-729.
- (76) Nishikawa M, Miyazaki C, Yamashita F, Takakura Y and Hashida M (1995) Galactosylated proteins are recognized by the liver according to the surface density of
20 galactose moieties. Am J Physiol 268, G849-856.

(77) Ong GL, Ettenson D, Sharkey RM, Marks A, Bauml R, Goldenberg DM and Mattes MJ

(1991) Galactose-conjugated antibodies in cancer therapy: properties and principles of action. Cancer Res 51, 1619-1626.

(78) Paganelli G, Stella M, De Nardi P, Magnani P, Zito F, Siccardi AG, Di Carlo V and

Fazio F (1991) A new method for faster blood clearance in radioimmuno-guided surgery.

J Nucl Biol Med 35, 88-89.

(79) Pai LH, FitzGerald DJ, Tepper M, Schacter B, Spitalny G and Pastan I (1990) Inhibition

of antibody response to Pseudomonas exotoxin and an immunotoxin containing

Pseudomonas exotoxin by 15-deoxyspergualin in mice. Cancer-Res 50, 7750-7753.

(80) Pedley RB, Boden JA, Boden R, Begent RH, Turner A, Haines AM and King DJ (1994)

The potential for enhanced tumour localisation by poly(ethylene glycol) modification of anti-CEA antibody. Br J Cancer 70, 1126-1130.

(81) Pedley RB, Dale R, Boden JA, Begent RH, Keep PA and Green AJ (1989) The effect of

second antibody clearance on the distribution and dosimetry of radiolabelled anti-CEA

antibody in a human colonic tumor xenograft model. Int J Cancer 43, 713-718.

(82) Pendri A, Conover CD and Greenwald RB (1998) Antitumor activity of paclitaxel-2'-

glycinate conjugated to poly(ethylene glycol): a water-soluble prodrug. Anticancer-Drug-Des 13, 387-395.

- (83) Rice KG, Weisz OA, Barthel T, Lee RT and Lee YC (1990) Defined geometry of binding between triantennary glycopeptide and the asialoglycoprotein receptor of rat hepatocytes. J Biol Chem 265, 18429-18434.
- (84) Richter AW and Akerblom E (1983) Antibodies against polyethylene glycol produced in animals by immunization with monomethoxy polyethylene glycol modified proteins. Int Arch Allergy Appl Immunol 70, 124-131.
- (85) Roffler SR, Chan J and Yeh MY (1994) Potentiation of radioimmunotherapy by inhibition of topoisomerase I. Cancer Res 54, 1276-1285.
- (86) Roffler SR, Wang SM, Chern JW, Yeh MY and Tung E (1991) Anti-neoplastic glucuronide prodrug treatment of human tumor cells targeted with a monoclonal antibody-enzyme conjugate. Biochem Pharmacol 42, 2062-2065.
- (87) Rogers GT, Burke PJ, Sharma SK, Koodie R and Boden JA (1995) Plasma clearance of an antibody-enzyme conjugate in ADEPT by monoclonal anti-enzyme: its effect on prodrug activation in vivo. Br J Cancer 72, 1357-1363.
- (88) Roseng L, Tolleshaug H and Berg T (1992) Uptake, intracellular transport, and degradation of polyethylene glycol-modified asialofetuin in hepatocytes. J-Biol-Chem 267, 22987-22993.
- (89) Ruddy SB and Hadzija BW (1994) High-performance liquid chromatographic method for the simultaneous determination of low-molecular-mass oligomers of polyethylene glycol in aqueous skin extracts. J-Chromatogr-B-Biomed-Appl 657, 83-92.

- (90) Ryan CM, Yarmush ML and Tompkins RG (1992) Separation and quantitation of polyethylene glycols 400 and 3350 from human urine by high-performance liquid chromatography. J-Pharm-Sci 81, 350-352.
- 5 (91) Saito T, Nishimura H, Sekine T, Urushibara T, Kodera Y, Hiroto M, Matsushima A and Inada Y (1996) Tolerogenic capacity of poly(ethylene glycol) (PEG)--modified ovalbumins in relation to their immunoreactivity towards anti-ovalbumin antibody. J Biomater Sci Polym Ed 8, 311-321.
- 10 (92) Sandborn WJ and Hanauer SB (1999) Antitumor necrosis factor therapy for inflammatory bowel disease: a review of agents, pharmacology, clinical results, and safety. Inflamm-Bowel-Dis 5, 119-133.
- (93) Savoca KV, Abuchowski A, van Es T, Davis FF and Palczuk NC (1979) Preparation of a non-immunogenic arginase by the covalent attachment of polyethylene glycol. Biochim-Biophys-Acta 578, 47-53.
- 15 (94) Savoca KV, Davis FF and Palczuk NC (1984) Induction of tolerance in mice by uricase and monomethoxypolyethylene glycol-modified uricase. Int Arch Allergy Appl Immunol 75, 58-67.
- (95) Senter PD, Saulnier MG, Schreiber GJ, Hirschberg DL, Brown JP, Hellstrom I and Hellstrom KE (1988) Anti-tumor effects of antibody-alkaline phosphatase conjugates in combination with etoposide phosphate. Proc Natl Acad Sci U S A 85, 4842-4846.

(96) Sharkey RM, Boerman OC, Natale A, Pawlyk D, Monestier M, Losman MJ and Goldenberg DM (1992) Enhanced clearance of radiolabeled murine monoclonal antibody by a syngeneic anti-idiotypic antibody in tumor-bearing nude mice. Int J Cancer 51, 266-273.

5 (97) Sharkey RM, Primus FJ and Goldenberg DM (1984) Second antibody clearance of radiolabeled antibody in cancer radioimmunodetection. Proc Natl Acad Sci U S A 81, 2843-2846.

(98) Sharma SK, Bagshawe KD, Burke PJ, Boden JA, Rogers GT, Springer CJ, Melton RG and Sherwood RF (1994) Galactosylated antibodies and antibody-enzyme conjugates in
10 antibody-directed enzyme prodrug therapy. Cancer 73, 1114-1120.

(99) Sharma SK, Bagshawe KD, Burke PJ, Boden RW and Rogers GT (1990) Inactivation and clearance of an anti-CEA carboxypeptidase G2 conjugate in blood after localisation in a xenograft model. Br J Cancer 61, 659-662.

(100) Sharma SK, Bagshawe KD, Melton RG and Begent RH (1996) Effect of cyclosporine on
15 immunogenicity of a bacterial enzyme carboxypeptidase G2 in ADEPT. Transplant Proc 28, 3154-3155.

(101) Stella M, De Nardi P, Paganelli G, Magnani P, Mangili F, Sassi I, Baratti D, Gini P, Zito F, Cristallo M and et al. (1994) Avidin-biotin system in radioimmunoguided surgery for colorectal cancer. Advantages and limits. Dis Colon Rectum 37, 335-343.

(102) Stewart JS, Sivolapenko GB, Hird V, Davies KA, Walport M, Ritter MA and Epenetos AA (1990) Clearance of ¹³¹I-labeled murine monoclonal antibody from patients' blood by intravenous human anti-murine immunoglobulin antibody. Cancer Res 50, 563-567.

5 (103) Stocks SJ, Jones AJ, Ramey CW and Brooks DE (1986) A fluorometric assay of the degree of modification of protein primary amines with polyethylene glycol. Anal Biochem 154, 232-234.

(104) Suzuki T, Kanbara N, Tomono T, Hayashi N and Shinohara I (1984) Physicochemical and biological properties of poly(ethylene glycol)-coupled immunoglobulin G. Biochim-Biophys-Acta 788, 248-255.

10 (105) Svensson HP, Frank IS, Berry KK and Senter PD (1998) Therapeutic effects of monoclonal antibody-beta-lactamase conjugates in combination with a nitrogen mustard anticancer prodrug in models of human renal cell carcinoma. J Med Chem 41, 1507-1512.

15 (106) Tennvall J, Garkavij M, Chen J, Sjogren HO and Strand SE (1997) Improving tumor-to-normal-tissue ratios of antibodies by extracorporeal immunoadsorption based on the avidin-biotin concept: development of a new treatment strategy applied to monoclonal antibodies murine L6 and chimeric BR96. Cancer 80, 2411-2418.

(107) Thor A, Ohuchi N, Szpak CA, Johnston WW and Schlom J (1986) Distribution of oncofetal antigen tumor-associated glycoprotein-72 defined by monoclonal antibody B72.3. Cancer Res 46, 3118-3124.

20

(108) Thornburg RW, Day JF, Baynes JW and Thorpe SR (1980) Carbohydrate-mediated clearance of immune complexes from the circulation. A role for galactose residues in the hepatic uptake of IgG-antigen complexes. J Biol Chem 255, 6820-6825.

5 (109) Townsend RR, Hardy MR, Wong TC and Lee YC (1986) Binding of N-linked bovine fetuin glycopeptides to isolated rabbit hepatocytes: Gal/GalNAc hepatic lectin discrimination between Gal beta(1,4)GlcNAc and Gal beta(1,3)GlcNAc in a triantennary structure. Biochemistry 25, 5716-5725.

10 (110) Ullen A, Riklund Ahlstrom K, Makiya R and Stigbrand T (1995a) Syngeneic anti-idiotypic antibodies eliminate excess radiolabeled idiotypes at experimental radioimmunolocalization. Cell Biophys 27, 31-45.

(111) Ullen A, Sandstrom P, Ahlstrom KR, Sundstrom B, Nilsson B, Arlestig L and Stigbrand T (1995b) Use of anticytokeratin monoclonal anti-idiotypic antibodies to improve tumor:nontumor ratio in experimental radioimmunolocalization. Cancer Res 55, 5868s-5873s.

15 (112) Vrudhula VM, Senter PD, Fischer KJ and Wallace PM (1993) Prodrugs of doxorubicin and melphalan and their activation by a monoclonal antibody-penicillin-G amidase conjugate. J Med Chem 36, 919-923.

20 (113) Wallace PM, MacMaster JF, Smith VF, Kerr DE, Senter PD and Cosand WL (1994) Intratumoral generation of 5-fluorouracil mediated by an antibody-cytosine deaminase conjugate in combination with 5-fluorocytosine. Cancer Res 54, 2719-2723.

(114) Wang QC, Pai LH, Debinski W, FitzGerald DJ and Pastan I (1993) Polyethylene glycol-modified chimeric toxin composed of transforming growth factor alpha and Pseudomonas exotoxin. Cancer-Res 53, 4588-4594.

5 (115) Wang SM, Chern JW, Yeh MY, Ng JC, Tung E and Roffler SR (1992) Specific activation of glucuronide prodrugs by antibody-targeted enzyme conjugates for cancer therapy. Cancer Res 52, 4484-4491.

(116) Yabe Y, Nishikawa M, Tamada A, Takakura Y and Hashida M (1999) Targeted delivery and improved therapeutic potential of catalase by chemical modification: combination with superoxide dismutase derivatives. J-Pharmacol-Exp-Ther 289, 1176-1184.

10 (117) Yeh MY, Hellstrom I, Brown JP, Warner GA, Hansen JA and Hellstrom KE (1979) Cell surface antigens of human melanoma identified by monoclonal antibody. Proc Natl Acad Sci U S A 76, 2927-2931.

15 The invention is not limited by the embodiments described above which are presented as examples only but can be modified in various ways within the scope of protection defined by the appended patent claims.